

## Product datasheet for **TP312844L**

### RASSF7 (NM\_003475) Human Recombinant Protein

#### Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human Ras association (RalGDS/AF-6) domain family (N-terminal) member 7 (RASSF7), transcript variant 1, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC212844 representing NM_003475 <b>Red</b> =Cloning site <b>Green</b> =Tags(s)
	<p>MLLGLAAMELKVWVDGIQRVVCVSEQTTCQEWIALAQAIGQTGRFVLVQRLREKERQLLPQCEPVGAQ ATCGQFASDVQFVLRRTGPSLAGRPSSDSCPPPERCLIRASLPVKPRAALGCEPRKTLTPEPAPLSRPG PAAPVTPTPGCCTDLRGLRLVQRNAEELGHEAFWEQELRREQAREREGQARLQALSAATAEHAARLQAL DAQARALEAELQLAAEAPGPPSPMASATERLHQDLAVQERQSAEVQGSALVSRALAAERALQAQAQEL EELNRELQCNLQQFIQQTGAALPPPPRPDRGPPGTQGPLPPAREESLLGAPSESHAGAQRPRRGPHDA ELLEVAAPAPEWCPLAAQPQAL</p> <p><b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b></p>
Tag:	C-Myc/DDK
Predicted MW:	39.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u><a href="#">NP_003466</a></u>



[View online >](#)

Locus ID:	8045
UniProt ID:	<a href="#">Q02833</a> , <a href="#">A0A024RCE4</a>
RefSeq Size:	1539
Cytogenetics:	11p15.5
RefSeq ORF:	1119
Synonyms:	C11orf13; CFAP88; FAP88; HRAS1; HRC1
Summary:	Negatively regulates stress-induced JNK activation and apoptosis by promoting MAP2K7 phosphorylation and inhibiting its ability to activate JNK. Following prolonged stress, anti-apoptotic effect stops because of degradation of RASSF7 protein via the ubiquitin-proteasome pathway. Required for the activation of AURKB and chromosomal congression during mitosis where it stimulates microtubule polymerization.[UniProtKB/Swiss-Prot Function]

### Product images:



Coomassie blue staining of purified RASSF7 protein (Cat# [TP312844]). The protein was produced from HEK293T cells transfected with RASSF7 cDNA clone (Cat# [RC212844]) using MegaTran 2.0 (Cat# [TT210002]).