

Product datasheet for **TP313056L**

Factor XI (F11) (NM_000128) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human coagulation factor XI (F11), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC213056 representing NM_000128 Red =Cloning site Green =Tags(s)

MIFLYQVVHFILFTSVSGECVTQLLKDTCFEGGDITTVFTPSAKYCQWCTYHPRCLLFTFTAESPSEDPTRWFTCVLKDSVTETLPRVNRATAISGYFSKQCSHQISACNKDIYVDLDMKGINYNSSVAKSAQECQERC
TDDVHCHFFTYATRQFPSLEHRNICLLKHTQTGTPTRITKLDKVVSGFSLKSCALSNLACIRDIFPNTVF
ADSNIDSVMAPDAFVCGRICTHHPGCLFFTFSSQEWPKESQRNLCLLKTSEGLPSTRIKKSALSGFSL
QSCRHSIPVFCHSSFYHDTDFLGEELDIVAAKSHEACQKLCTNAVRCQFFTYTPAQASCNEGKGKCYLKL
SSNGSPTKILHGRGGISGYTLRLCKMDNECTTKIKPRIVGGTASVRGEWPWQVTLHTTSPTQRHLCCGSI
IGNQWILTAAHCFYGVESPKILRVYSGILNQSEIKEDTSFFGVQEIIHDQYKMAESGYDIALLKLETTV
NYTDSQRPICLPSKGDRNVIYTDWVTGWGYRKLQNTLQKAKIPLVTNEECQKRYRGHKITHKMIC
AGYREGGKDACKGDSGGPLSCKHNEVWHLVGITSWGEGCAQRERPGVYTNWEYVDWILEKTQAV

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	69.9 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_000119</u>
Locus ID:	2160
UniProt ID:	<u>P03951</u>
RefSeq Size:	2217
Cytogenetics:	4q35.2
RefSeq ORF:	1875
Synonyms:	FXI; PTA
Summary:	This gene encodes coagulation factor XI of the blood coagulation cascade. This protein is present in plasma as a zymogen, which is a unique plasma coagulation enzyme because it exists as a homodimer consisting of two identical polypeptide chains linked by disulfide bonds. During activation of the plasma factor XI, an internal peptide bond is cleaved by factor XIIa (or XII) in each of the two chains, resulting in activated factor XIa, a serine protease composed of two heavy and two light chains held together by disulfide bonds. This activated plasma factor XI triggers the middle phase of the intrinsic pathway of blood coagulation by activating factor IX. Defects in this factor lead to Rosenthal syndrome, a blood coagulation abnormality. [provided by RefSeq, Jul 2008]
Protein Families:	Druggable Genome, Protease, Secreted Protein
Protein Pathways:	Complement and coagulation cascades

Product images:



Coomassie blue staining of purified F11 protein (Cat# [TP313056]). The protein was produced from HEK293T cells transfected with F11 cDNA clone (Cat# [RC213056]) using MegaTran 2.0 (Cat# [TT210002]).