

Product datasheet for **TP313140**

WFIKKN2 (NM_175575) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human WAP, follistatin/kazal, immunoglobulin, kunitz and netrin domain containing 2 (WFIKKN2), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA	>RC213140 representing NM_175575
Clone or AA Sequence:	Red=Cloning site Green=Tags(s)

MWAPRCRRFWSRWEQVAALLLLLLLLLVPPRSLALPPIRYSHAGICPNDMNPNLWVDAQSTCRRECETDQ
ECETYEKCCPNVCGTKSCVAARYMDVKGKKGVPVGMPEATCDHFMCLQQGSECDIWDGQPVCKCKDRCEK
EPSFTCASDGLTYNRCYMDAEACSKGITLAWVTCRYHFTWPNTSPPPPETTMHPTTASPETPELDMAAP
ALLNNPVHQSVTMGETVSFLCDVWGRPRPEITWEKQLEDRENVVMRPNHVRGNVVVTNIAQLVIYNAQLQ
DAGIYTCTARNVAGVLRADFPLSVVRGHQAAATSESSPNGTAFPAECLKPPDSEDCGEEQTRWHFDAQA
NNCLTFTFGHCHRNLNHFETYEACMLACMSGPLAACSLPALQGPKAYAPRWAYNSQTGQCQSFVYGGCE
GNGNNFESREACEESCPFPRGNQRCRACKPRQLVTSFCRSDFVILGRVSELTEEPDSGRALVTVDEVLK
DEKMGLKFLGQEPLEVTLLHVDWACPCPNVTSEMPLIIMGEVDGGMAMLRPDSFVGASSARRVRKLREV
MHKKTCDVLKEFLGLH

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	63.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_783165
Locus ID:	124857
UniProt ID:	Q8TEU8
RefSeq Size:	3534
Cytogenetics:	17q21.33
RefSeq ORF:	1728
Synonyms:	GASP-1; hGASP-1; WFDC20B; WFIKKNRP
Summary:	The WFIKKN1 protein contains a WAP domain, follistatin domain, immunoglobulin domain, two tandem Kunitz domains, and an NTR domain. This gene encodes a WFIKKN1-related protein which has the same domain organization as the WFIKKN1 protein. The WAP-type, follistatin type, Kunitz-type, and NTR-type protease inhibitory domains may control the action of multiple types of proteases. [provided by RefSeq, Jul 2008]
Protein Families:	Transmembrane

Product images:



Coomassie blue staining of purified WFIKKN2 protein (Cat# TP313140). The protein was produced from HEK293T cells transfected with WFIKKN2 cDNA clone (Cat# [RC213140]) using MegaTran 2.0 (Cat# [TT210002]).