

## Product datasheet for TP313298

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## APE1 (APEX1) (NM 080649) Human Recombinant Protein

**Product data:** 

**Product Type: Recombinant Proteins** 

Recombinant protein of human APEX nuclease (multifunctional DNA repair enzyme) 1 (APEX1), Description:

transcript variant 3, 20 µg

Species: Human **Expression Host:** HEK293T

Expression cDNA Clone >RC213298 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

> MPKRGKKGAVAEDGDELRTEPEAKKSKTAAKKNDKEAAGEGPALYEDPPDQKTSPSGKPATLKICSWNVD GLRAWIKKKGLDWVKEEAPDILCLQETKCSENKLPAELQELPGLSHQYWSAPSDKEGYSGVGLLSRQCPL KVSYGIGDEEHDQEGRVIVAEFDSFVLVTAYVPNAGRGLVRLEYRQRWDEAFRKFLKGLASRKPLVLCGD LNVAHEEIDLRNPKGNKKNAGFTPQERQGFGELLQAVPLADSFRHLYPNTPYAYTFWTYMMNARSKNVGW

RLDYFLLSHSLLPALCDSKIRSKALGSDHCPITLYLAL

**TRTRPLEOKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK Predicted MW:

Concentration: >0.05 µg/µL as determined by microplate BCA method

35.4 kDa

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Store at -80°C. Storage:

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 542380

Locus ID: 328





UniProt ID: P27695

RefSeq Size: 1507

Cytogenetics: 14q11.2

RefSeq ORF: 954

Synonyms: APE; APE1; APEN; APEX; APX; HAP1; REF1

**Summary:** The APEX gene encodes the major AP endonuclease in human cells. It encodes the APEX

> endonuclease, a DNA repair enzyme with apurinic/apyrimidinic (AP) activity. Such AP activity sites occur frequently in DNA molecules by spontaneous hydrolysis, by DNA damaging agents or by DNA glycosylases that remove specific abnormal bases. The AP sites are the most frequent pre-mutagenic lesions that can prevent normal DNA replication. Splice variants have been found for this gene; all encode the same protein. Disruptions in the biological functions


related to APEX are associated with many various malignancies and neurodegenerative

diseases.[provided by RefSeq, Dec 2019]

**Protein Families:** Druggable Genome, Stem cell - Pluripotency, Transcription Factors

**Protein Pathways:** Base excision repair

## **Product images:**



Coomassie blue staining of purified APEX1 protein (Cat# TP313298). The protein was produced from HEK293T cells transfected with APEX1 cDNA clone (Cat# [RC213298]) using

MegaTran 2.0 (Cat# [TT210002]).