

Product datasheet for TP313722

SLC39A7 (NM_001077516) Human Recombinant Protein

Product data:

| | |
|---------------------------------------|--|
| Product Type: | Recombinant Proteins |
| Description: | Recombinant protein of human solute carrier family 39 (zinc transporter), member 7 (SLC39A7), transcript variant 2, 20 µg |
| Species: | Human |
| Expression Host: | HEK293T |
| Expression cDNA Clone or AA Sequence: | >RC213722 protein sequence Red=Cloning site Green=Tags(s) MARGLGAPHWVAVGLLTWATLGLLVAGLGGHDDLHDDLQEDFHGHSRHSHEDFHHGHSHAHGHGHTHES IWHGHTHDHDHGHSHEDLHHGHSHGYSHESLYHRGHGHDHEHSHGGYGESGAPGIKQDLDAVTLWAYALG ATVLISAAPFFVFLIPVESNSPRHRSLLQILLSFASGGLLDGDAFLHLIPHAPHSHHTLEQPGHGHSH SGQGPILSVGLWVLSGIVAFVVEKFRVHVKGHGHSHGHGHAHSHTRGSHGHGRQERSTKEKQSSEEEE KETRGVQKRRGGSTVPKDGVPVPQNAEEERGLDLRVSGYLNLAADLAHNFTDGLAIGASFRGGRGLGIL TTMTVLLHEVPHEVGDFAILVQSGCSKKQAMRLQLLTAVGALAGTACALLTEGGAVGSEIAGGAGPGWVL PFTAGGFIYVATVSVLPELLREASPLQSLLEVLGLLGGVIMMVLIAHLE TRTRPLEQKLISEEDLAANDILDYKDDDDKV |
| Tag: | C-Myc/DDK |
| Predicted MW: | 49.9 kDa |
| Concentration: | >0.05 µg/µL as determined by microplate BCA method |
| Purity: | > 80% as determined by SDS-PAGE and Coomassie blue staining |
| Buffer: | 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol |
| Preparation: | Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps. |
| Note: | For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process. |
| Storage: | Store at -80°C. |
| Stability: | Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles. |



[View online »](#)

RefSeq: [NP_001070984](#)

Locus ID: 7922

UniProt ID: [Q92504](#), [A0A024RCX7](#)

RefSeq Size: 2172

Cytogenetics: 6p21.32


RefSeq ORF: 1407

Synonyms: D6S115E; D6S2244E; H2-KE4; HKE4; KE4; RING5; ZIP7

Summary: The protein encoded by this gene transports zinc from the Golgi and endoplasmic reticulum to the cytoplasm. This transport may be important for activation of tyrosine kinases, some of which could be involved in cancer progression. Therefore, modulation of the encoded protein could be useful as a therapeutic agent against cancer. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2014]

Protein Families: Transmembrane

Product images:



Coomassie blue staining of purified SLC39A7 protein (Cat# TP313722). The protein was produced from HEK293T cells transfected with SLC39A7 cDNA clone (Cat# [RC213722]) using MegaTran 2.0 (Cat# [TT210002]).