

Product datasheet for TP313770

FBXO40 (NM_016298) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human F-box protein 40 (FBXO40), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC213770 representing NM_016298 Red=Cloning site Green=Tags(s)

MGKARRSPPGHHRHCEGCFNRHCHIPVEPNTSCLVISCHLLCGATFHMCKEAEHQLLCPLEQVPCLNSEY
GCPLSMSRHKLAKHLQVCPASVCCSMEWNRWPNVDSETTLHENIMKETPSEECLDTALALQDQKVLFRS
LKMVELFPETREATEEEPTMNGETSVEEMGGAVGGVDIGLVPGLSATNGEMAELSQEEREVLAKTKEGM
DLVKGQWENIFSKEHAASALTNSSASCESKNKNDSEKEQISSGHNMVEGEGAPKKKEPQENKQKQDVRT
AMETTGLAPWQDGVLERLKTAVDAKDYNMYLVHNGRMLIHFGQMPACTPKERDFVYGKLEAQEVKTVYTF
KVPVSYCGKRARLGDAMLSCKPSEHKAVDTSDLGITVEDLPKSDLIKTTLQCALERELKGHVISESRSID
GLFMDFATQTYNFEPEQFSSGTVLADLTAATPGGLHVELHSECVTRRHNKSSSAFTFTCNKFFRRDEFPL
HFKNVHTDIQSCLNGWFQHRCPPLAYLGCTFVQNHFRPPGQKAKVIYSQELKTFAIKPEVAPELSEGRKNN
HLLGHGGKSQNSLTSLPLEILKYIAGFLDSVSLAQLSQVSVLMRNICATLLQERGMVLLQWKKKRYSHGG
TSWRVHREIWQFSSLFSKIKSWEFNEVTSMSSEHLKSCPFNIVEHKTDPILLTSMCQPREQARESLVSTFR
IRPRGRYVS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	79.6 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_057382](#)

Locus ID: 51725

UniProt ID: [Q9UH90](#)

RefSeq Size: 3442

Cytogenetics: 3q13.33


RefSeq ORF: 2127

Synonyms: FBX40

Summary: Members of the F-box protein family, such as FBXO40, are characterized by an approximately 40-amino acid F-box motif. SCF complexes, formed by SKP1 (MIM 601434), cullin (see CUL1; MIM 603134), and F-box proteins, act as protein-ubiquitin ligases. F-box proteins interact with SKP1 through the F box, and they interact with ubiquitination targets through other protein interaction domains (Jin et al., 2004 [PubMed 15520277]).[supplied by OMIM, Mar 2008]

Protein Families: Druggable Genome

Product images:



Coomassie blue staining of purified FBXO40 protein (Cat# TP313770). The protein was produced from HEK293T cells transfected with FBXO40 cDNA clone (Cat# [RC213770]) using MegaTran 2.0 (Cat# [TT210002]).