

## Product datasheet for TP313845

### CD96 (NM\_198196) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human CD96 molecule (CD96), transcript variant 1, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC213845 representing NM_198196 Red=Cloning site Green=Tags(s)

MEKKWKYCAVYYIIQIHFKGVWEKTVNTEENVYATLGSDVNLTCQTQTVGFFVQMQWSKVTNKIDLIAV  
YHPQYGFYCAGRPCESLVTFTETPENGSKWTLHLRNMSCSVSGRYECMLVLYPEGIQTKIYNLLIQTHV  
TADEWNSNHTIEIEINQTL EIPCFQNSSKISSEFTYAWSVENSSTDSWLLSKGIKEDNGTQETLISQN  
HLISNSTLLKDRVKLGTDYRLHLSVPVQIFDDGRKFSCHIRVGNPKILRSSTTVKVFAPKEIPVIVENNST  
DVLVERRFTCLLKNVFPKANITW FIDG SFLHDEKEGIYITNEERKKGKDGFLKSVLTRVHSNKPAQSDN  
LTIWCMALSPVPGNKVWNISSEKITFLLGSEISSTDPPLSVTESTLDTQPSPASSVSPARYPATSSVTLV  
DVSALRPNTTPQPSNSSMTTRGFNYPWTSSGTDTKKSVSRIPSETYSSSPSGAGSTLHDNVFTSTARAFS  
EVPTTANGSTKTNHVHITGIVVNKPKDGMSPVIVAALLFCCMILFGLGVRKWCQYQKEIMERPPPFKPP  
PPPIKYTCIQEPNESDLPYHEMETL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	63 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<a href="#">NP_937839</a>
<b>Locus ID:</b>	10225
<b>UniProt ID:</b>	<a href="#">P40200</a>
<b>RefSeq Size:</b>	4561
<b>Cytogenetics:</b>	3q13.13-q13.2
<b>RefSeq ORF:</b>	1755
<b>Synonyms:</b>	TACTILE
<b>Summary:</b>	The protein encoded by this gene belongs to the immunoglobulin superfamily. It is a type I membrane protein. The protein may play a role in the adhesive interactions of activated T and NK cells during the late phase of the immune response. It may also function in antigen presentation. Alternative splicing generates multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Jan 2016]
<b>Protein Families:</b>	Druggable Genome, Transmembrane

### Product images:



Coomassie blue staining of purified CD96 protein (Cat# TP313845). The protein was produced from HEK293T cells transfected with CD96 cDNA clone (Cat# [RC213845]) using MegaTran 2.0 (Cat# [TT210002]).