

## Product datasheet for TP313966M

### NFAT1 (NFATC2) (NM\_173091) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 2 (NFATC2), transcript variant 2, 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC213966 representing NM_173091 Red=Cloning site Green=Tags(s)

MNAPERQPQPDGGDAPGHEPGGSPQDELDFSI LFDY EYLNPN EEPNAHKVASPPSGPAYPDDVLDYGLK  
 PYSPLASLSGEPGRFGEPDRVGPQKFLSAAK PAGASGLSPRIEITPSHEL IQAVGPLRMRDAGLLVEQP  
 PLAGVAASPRFTLPVPGFEGYREPLCLSPASSGSSASFISDTFSPYTPCVSPNNGGPDLLCPQFQNI PA  
 HYSPTSPIMSPRTSLAEDSCLGRHSPVPRPASRSSSPGAKRRHS CAEALVALPPGASQRRSRSPSPQPS  
 SHVAPQDHGSPAGYPPVAGSAVIMDALNSLATDSPCGIPPKMWKTS PDPSPVSAAPSKAGLPRHIYPAVE  
 FLGPCEQGERRNSAPESILLVPPTWPKPLVPAIPIC SIPVTASLPLEWPLSSQSGSYELRIEVQPKPHH  
 RAHYETEGSRGAVKAPTGGHPVQVLHGYMENKPLGLQIFIGTAD ERILKPHAFYQVHRITGKTVTTTSYE  
 KIVGNTKVL EIPLEPKNNMRATIDCAGILKLRNADIELRKG ETDIGRKNTRVRLVFRVHIPESSGRIVSL  
 QTASNPIECSQRSAHELPMVERQD TDSCLVYGGQQMIL TGQNFTSESKVVFTEKTTDGGQI WEMEATVDK  
 DKSQPNMLFVEIPEYRNKHIRTPVKVNFYVINGKRKRSQPQHFTYHPVPAIKTEPTDEYDPTLICSP THG  
 GLGSQPYYPQHMPVAESPSCLVATMAPCQQFRTGLSSPDARYQQQNPAAVLYQRSKSLSPSLLGYQQPAL  
 MAAPLSLADAHRSVLVHAGSQGQSSALLHPSPTNQQASPV IHYSPTNQQLRGCGSHQEFQHIMYCENFAPG  
 TTRPGPPPVSQGRQLSPGSYPTVIQQNATSQRAAKNGPPVSDQKEVLPAGVTIKQEQLNDQTYLDDVNE  
 IIRKEFSGPPARNQT

TRTRPLEQKLI SEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	100 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol



[View online »](#)

<b>Preparation:</b>	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
<b>Note:</b>	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
<b>Storage:</b>	Store at -80°C.
<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<a href="#">NP_775114</a>
<b>Locus ID:</b>	4773
<b>UniProt ID:</b>	<a href="#">Q13469</a>
<b>RefSeq Size:</b>	3166
<b>Cytogenetics:</b>	20q13.2
<b>RefSeq ORF:</b>	2775
<b>Synonyms:</b>	NFAT1; NFATP
<b>Summary:</b>	This gene is a member of the nuclear factor of activated T cells (NFAT) family. The product of this gene is a DNA-binding protein with a REL-homology region (RHR) and an NFAT-homology region (NHR). This protein is present in the cytosol and only translocates to the nucleus upon T cell receptor (TCR) stimulation, where it becomes a member of the nuclear factors of activated T cells transcription complex. This complex plays a central role in inducing gene transcription during the immune response. Alternate transcriptional splice variants encoding different isoforms have been characterized. [provided by RefSeq, Apr 2012]
<b>Protein Families:</b>	Druggable Genome, Transcription Factors
<b>Protein Pathways:</b>	Axon guidance, B cell receptor signaling pathway, MAPK signaling pathway, Natural killer cell mediated cytotoxicity, T cell receptor signaling pathway, VEGF signaling pathway, Wnt signaling pathway

### Product images:



Coomassie blue staining of purified NFATC2 protein (Cat# [TP313966]). The protein was produced from HEK293T cells transfected with NFATC2 cDNA clone (Cat# [RC213966]) using MegaTran 2.0 (Cat# [TT210002]).