

## Product datasheet for TP314065L

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## p19 INK4d (CDKN2D) (NM 001800) Human Recombinant Protein

**Product data:** 

**Product Type: Recombinant Proteins** 

Recombinant protein of human cyclin-dependent kinase inhibitor 2D (p19, inhibits CDK4) Description:

(CDKN2D), transcript variant 1, 1 mg

Species: Human **Expression Host:** HEK293T

**Expression cDNA Clone** >RC214065 representing NM 001800 or AA Sequence:

Red=Cloning site Green=Tags(s)

MLLEEVRAGDRLSGAAARGDVQEVRRLLHRELVHPDALNRFGKTALQVMMFGSTAIALELLKQGASPNVQ DTSGTSPVHDAARTGFLDTLKVLVEHGADVNVPDGTGALPIHLAVQEGHTAVVSFLAAESDLHRRDARGL

**TPLELALQRGAQDLVDILQGHMVAPL** 

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK Predicted MW: 17.5 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

> 80% as determined by SDS-PAGE and Coomassie blue staining **Purity:** 

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Recombinant protein was captured through anti-DDK affinity column followed by conventional **Preparation:** 

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Store at -80°C. Storage:

Stable for 12 months from the date of receipt of the product under proper storage and Stability:

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001791

1032 Locus ID:

UniProt ID: P55273, A0A024R796





RefSeq Size: 1416

Cytogenetics: 19p13.2 RefSeq ORF: 498

Synonyms: INK4D; p19; p19-INK4D

Summary: The protein encoded by this gene is a member of the INK4 family of cyclin-dependent kinase

> inhibitors. This protein has been shown to form a stable complex with CDK4 or CDK6, and prevent the activation of the CDK kinases, thus function as a cell growth regulator that controls cell cycle G1 progression. The abundance of the transcript of this gene was found to oscillate in a cell-cycle dependent manner with the lowest expression at mid G1 and a maximal expression during S phase. The negative regulation of the cell cycle involved in this

protein was shown to participate in repressing neuronal proliferation, as well as


spermatogenesis. Two alternatively spliced variants of this gene, which encode an identical

protein, have been reported. [provided by RefSeq, Jul 2008]

**Protein Families:** Druggable Genome

**Protein Pathways:** Cell cycle

## **Product images:**



Coomassie blue staining of purified CDKN2D protein (Cat# [TP314065]). The protein was produced from HEK293T cells transfected with CDKN2D cDNA clone (Cat# [RC214065]) using

MegaTran 2.0 (Cat# [TT210002]).