

Product datasheet for **TP314079L**

Zyxin (ZYG) (NM_003461) Human Recombinant Protein

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human zyxin (ZYX), transcript variant 1, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC214079 representing NM_003461 Red =Cloning site Green =Tags(s)
	MAAPRPSPAISVSVSAPAFYAPQKKFGPVVAPKPKVNPFRPGDSEPPAPGAQRAQMGRVGEIPPPPPED FPLPPPPLAGDGDDAEGALGGAFPPPPPIEESFPPAPLEEEIFSPPPPPPEEGGPEAPIPPPPQPREK VSSIDLEIDSLSSLLDDMTKNDPFKARVSSGYVPPPVPATPFSSKSSTKPAAGGTAPLPPWKSPSSSQPLP QVPAPAQSQTQFHVQPQPQPKPQVQLHVQSQTQPVSLANTQPRGPPASSPAPAPKFSVPTPKFTPVASKF SPGAPGGSGSQPNQKLGHPALSAGTGSPQPPSFTYAQQREKPRVQEKQHPVPPPAQNQNQVRSFGAPGP LTLKEVEELEQLTQQLMQDMEHPQRQNVAVNELCGRCHQPLARAQPAVRALGQLFHIACTCHQCAQQLQ GQQFYSLEGAPYCEGCYTDLTLEKNTCGEPITDRMLRATGKAYHPHCFTCVCARPLEGTSFIVDQANRP HCVDPDYHKQYAPRCSVCSEPIMPEPGRDETVRVVALDKNFHMKCYKCEDCGKPLSIEADDNGCFPLDGHV LCRKCHTARAQT
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	61.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_003452
Locus ID:	7791
UniProt ID:	Q15942 , Q96AF9
RefSeq Size:	2325
Cytogenetics:	7q34
RefSeq ORF:	1716
Synonyms:	ESP-2; HED-2
Summary:	Focal adhesions are actin-rich structures that enable cells to adhere to the extracellular matrix and at which protein complexes involved in signal transduction assemble. Zyxin is a zinc-binding phosphoprotein that concentrates at focal adhesions and along the actin cytoskeleton. Zyxin has an N-terminal proline-rich domain and three LIM domains in its C-terminal half. The proline-rich domain may interact with SH3 domains of proteins involved in signal transduction pathways while the LIM domains are likely involved in protein-protein binding. Zyxin may function as a messenger in the signal transduction pathway that mediates adhesion-stimulated changes in gene expression and may modulate the cytoskeletal organization of actin bundles. Alternative splicing results in multiple transcript variants that encode the same isoform. [provided by RefSeq, Jul 2008]
Protein Pathways:	Focal adhesion

Product images:



Coomassie blue staining of purified ZYX protein (Cat# [TP314079]). The protein was produced from HEK293T cells transfected with ZYX cDNA clone (Cat# [RC214079]) using MegaTran 2.0 (Cat# [TT210002]).