

Product datasheet for **TP314156L**

Bestrophin (BEST1) (NM_004183) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human bestrophin 1 (BEST1), transcript variant 1, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC214156 representing NM_004183 Red =Cloning site Green =Tags(s)

MTITYTSQVANARLGSFSRLLLCWRGSIYKLLYGEFLIFLLCYIIRFIYRLALTEEQQLMFEKLTLYCD
SYIQLIPISFVLGFYVTLVWTRWWNQYENLPWPDRLMSLVSGFVEGKDEQGRLLRRTLIRYANLGNVLIL
RSVSTAVYKRFPSAQHLVQAGFMTPAEHKQLEKLSLPHNMFVWPVWFANLSMKAWLGGRIKDPILLQSL
LNEMNLTLCQCGHLYAYDWISIPVYTVAVYSFFLTCLVGRQFLNPAKAYPGHELDLVVPVFTFLQ
FFFYVGVWLKVAEQLINPFGEDDDDFETNWIVDRNLQVSLAVDEMHDLPMEPDMYWNKPEPQPPYTAA
SAQFRRASFMGSTFNISLNKEEMEFQPNQEDEEDAHAGIIGRFLGLQSHDHHPPRANSRTKLLWPKRESL
LHEGLPKNHKAAKQNVRGQEDNKAWKLVKAVDAFKSAPLYQRPQGYSSAPQTPSPTPMFFPLEPSAPSKLH
SVTGIDTKDKSLKTVSSGAKKSFELLESSESGALMEHPEVSVRRKTVFNLDTMPEIPENHLKEPLEQSP
TNIHTTLKDHMDPYWALENRDEAHS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	67.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_004174
Locus ID:	7439
UniProt ID:	O76090
RefSeq Size:	2673
Cytogenetics:	11q12.3
RefSeq ORF:	1755
Synonyms:	ARB; BEST; Best1V1Delta2; BMD; RP50; TU15B; VMD2
Summary:	This gene encodes a member of the bestrophin gene family. This small gene family is characterized by proteins with a highly conserved N-terminus with four to six transmembrane domains. Bestrophins may form chloride ion channels or may regulate voltage-gated L-type calcium-ion channels. Bestrophins are generally believed to form calcium-activated chloride-ion channels in epithelial cells but they have also been shown to be highly permeable to bicarbonate ion transport in retinal tissue. Mutations in this gene are responsible for juvenile-onset vitelliform macular dystrophy (VMD2), also known as Best macular dystrophy, in addition to adult-onset vitelliform macular dystrophy (AVMD) and other retinopathies. Alternative splicing results in multiple variants encoding distinct isoforms.[provided by RefSeq, Nov 2008]
Protein Families:	Druggable Genome, Ion Channels: Other, Transmembrane

Product images:



Coomassie blue staining of purified BEST1 protein (Cat# [TP314156]). The protein was produced from HEK293T cells transfected with BEST1 cDNA clone (Cat# [RC214156]) using MegaTran 2.0 (Cat# [TT210002]).