

Product datasheet for **TP314169M**

CLLD7 (RCBTB1) (NM_018191) Human Recombinant Protein

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human regulator of chromosome condensation (RCC1) and BTB (POZ) domain containing protein 1 (RCBTB1), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC214169 representing NM_018191 Red =Cloning site Green =Tags(s)
	<p>MVDVGKWPIFTLLSPQEIASIRKACVFGTSASEALYVTDNDEVFVGLNYSNCLGTGDNQSTLVPKKLEG LCGKKIKSLSYSGSPHVLLSTEDGVVYAWGHNGYSQLGNGTTNQGAPVQVCTNLLIKQVVEVACGSHHS MALAADGEVFAWGYNNCGQVSGSTANQPTPRKVTNCLHIKRVVGIACGQTSSMAVLDNGEVYWGYN NGN GQLGLGNNGNQLTTPVRVAALHSVCVNQIVCGYAHTLALTDEGLLYAWGANTYQGLGTGNKNLLSPA HIM VEKERVVEIAACHSAHTSAAKTQGGHVYMWGQCRGQSVILPHLTHFSCTDDVFACFATPAVSWRLLSVEH EDFLTVAESLKKEFDSPETADLKFRIDGKYIHVHKAVLKIRCEHFRSMFQSYWNEDMKEVIEIDQFSYPV YRAFLQYLYTDTVDLPPEDAIGLLDLATSYCENRLKKLCQHIIKRGITVENAFSLFSAAVRYDAEDLEEF CFKFCINHLTEVTQTAAFWQMDGPLLKEFIAKASKCGAFKN</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-Myc/DDK
Predicted MW:	58.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_060661</u>
Locus ID:	55213
UniProt ID:	<u>Q8NDN9</u>
RefSeq Size:	4042
Cytogenetics:	13q14.2
RefSeq ORF:	1593
Synonyms:	CLLD7; CLLL7; GLP; RDEOA
Summary:	This gene encodes a protein with an N-terminal RCC1 domain and a C-terminal BTB (broad complex, tramtrack and bric-a-brac) domain. In rat, over-expression of this gene in vascular smooth muscle cells induced cellular hypertrophy. In rat, the C-terminus of RCBTB1 interacts with the angiotensin II receptor-1A. In humans, this gene maps to a region of chromosome 13q that is frequently deleted in B-cell chronic lymphocytic leukemia and other lymphoid malignancies. [provided by RefSeq, Jul 2008]

Product images:



Coomassie blue staining of purified RCBTB1 protein (Cat# [TP314169]). The protein was produced from HEK293T cells transfected with RCBTB1 cDNA clone (Cat# [RC214169]) using MegaTran 2.0 (Cat# [TT210002]).