

# Product datasheet for TP314885M

# H2AC14 (NM\_021066) Human Recombinant Protein

# **Product data:**

#### **Product Type: Recombinant Proteins** Recombinant protein of human histone cluster 1, H2aj (HIST1H2AJ), 100 µg **Description:** Species: Human HEK293T **Expression Host:** Expression cDNA Clone >RC214885 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s) MSGRGKQGGKARAKAKTRSSRAGLQFPVGRVHRLLRKGNYAERVGAGAPVYLAAVLEYLTAEILELAGNA ARDNKKTRIIPRHLQLAIRNDEELNKLLGKVTIAQGGVLPNIQAVLLPKKTESHHKTK **TRTRPLEQKLISEEDLAANDILDYKDDDDKV** Tag: C-Myc/DDK Predicted MW: 13.8 kDa **Concentration:** >0.05 µg/µL as determined by microplate BCA method **Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining **Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol Recombinant protein was captured through anti-DDK affinity column followed by **Preparation:** conventional chromatography steps. Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process. Store at -80°C. Storage: Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles. RefSeq: NP 066544 Locus ID: 8331 **UniProt ID:** 099878 **RefSeq Size:** 439 Cytogenetics: 6p22.1



View online »


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	H2AC14 (NM_021066) Human Recombinant Protein – TP314885M
RefSeq ORF:	384
Synonyms:	dJ160A22.4; H2A/E; H2AFE; HIST1H2AJ
Summary:	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H2A family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Aug 2015]
Protein Pathway	s: Systemic lupus erythematosus

## **Product images:**



Coomassie blue staining of purified HIST1H2AJ protein (Cat# [TP314885]). The protein was produced from HEK293T cells transfected with HIST1H2AJ cDNA clone (Cat# [RC214885]) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US