

Product datasheet for **TP314987M**

TGF beta 3 (TGFB3) (NM_003239) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human transforming growth factor, beta 3 (TGFB3), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC214987 representing NM_003239 Red =Cloning site Green =Tags(s)

MKMHLQRALWVLLNLFATVLSLSTCTTDFGHIKKRVEAIRGQILSKLRLTSPPEPTVMTHVPYQVL
ALYNSTRELLEEMHGEREEGCTQENTESEYYAKEIHKFDMIQGLAEHNELAVCPKGITSKVFRFNVSVE
KNRTNLFRAEFRVLRVNPSSKRNEQRIELFQILRPDEHIAKQRYIGGKNLPTRGTAEWLSFDVTDVRE
WLLRRESNLGLEISIHCPCHTFQPNGDILENIHEVMEIKFKGVDNEDDHGRGDLGRLKKQKDHHNPHLIL
MMIPPHRLDNPQGQGRKKRALDTNYCFRNLEENCCVRPLYIDFRQDLGWKWWHEPKGYANFCSGPCPY
LRSADTTHSTVLGLYNTLNPEASASPCCVPQDLEPLTILYVGRTPKVEQLSNMVKSKCS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	44.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_003230
Locus ID:	7043



[View online »](#)

UniProt ID:	P10600 , A5YM40 , B3KVH9
RefSeq Size:	2574
Cytogenetics:	14q24.3
RefSeq ORF:	1236
Synonyms:	ARVD; ARVD1; LDS5; RNHF; TGF-beta3
Summary:	This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate a latency-associated peptide (LAP) and a mature peptide, and is found in either a latent form composed of a mature peptide homodimer, a LAP homodimer, and a latent TGF-beta binding protein, or in an active form consisting solely of the mature peptide homodimer. The mature peptide may also form heterodimers with other TGF-beta family members. This protein is involved in embryogenesis and cell differentiation, and may play a role in wound healing. Mutations in this gene are a cause of aortic aneurysms and dissections, as well as familial arrhythmogenic right ventricular dysplasia 1. [provided by RefSeq, Aug 2016]
Protein Families:	Druggable Genome, Secreted Protein, Transmembrane
Protein Pathways:	Cell cycle, Chronic myeloid leukemia, Colorectal cancer, Cytokine-cytokine receptor interaction, Dilated cardiomyopathy, Hypertrophic cardiomyopathy (HCM), MAPK signaling pathway, Pancreatic cancer, Pathways in cancer, Renal cell carcinoma, TGF-beta signaling pathway

Product images:



Coomassie blue staining of purified TGFB3 protein (Cat# [TP314987]). The protein was produced from HEK293T cells transfected with TGFB3 cDNA clone (Cat# [RC214987]) using MegaTran 2.0 (Cat# [TT210002]).