

Product datasheet for TP315248M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RAP1A (NM 002884) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human RAP1A, member of RAS oncogene family (RAP1A), transcript

variant 2, 100 µg

Species: Human Expression Host: HEK293T

Expression cDNA Clone >RC215248 representing NM_002884 **or AA Sequence:** Red=Cloning site Green=Tags(s)

MREYKLVVLGSGGVGKSALTVQFVQGIFVEKYDPTIEDSYRKQVEVDCQQCMLEILDTAGTEQFTAMRDL YMKNGQGFALVYSITAQSTFNDLQDLREQILRVKDTEDVPMILVGNKCDLEDERVVGKEQGQNLARQWCN

CAFLESSAKSKINVNEIFYDLVRQINRKTPVEKKKPKKKSCLLL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 20.8 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 002875

Locus ID: 5906

UniProt ID: <u>P62834</u>, <u>A8KAH9</u>





RefSeq Size: 1812

Cytogenetics: 1p13.2 RefSeq ORF: 552

Synonyms: C21KG; G-22K; KREV-1; KREV1; RAP1; SMGP21

Summary: This gene encodes a member of the Ras family of small GTPases. The encoded protein

undergoes a change in conformational state and activity, depending on whether it is bound to GTP or GDP. This protein is activated by several types of guanine nucleotide exchange factors (GEFs), and inactivated by two groups of GTPase-activating proteins (GAPs). The activation status of the encoded protein is therefore affected by the balance of intracellular levels of GEFs and GAPs. The encoded protein regulates signaling pathways that affect cell proliferation and adhesion, and may play a role in tumor malignancy. Pseudogenes of this gene have been defined on chromosomes 14 and 17. Alternative splicing results in multiple transcript variants.

[provided by RefSeq, May 2014]


Protein Families: Druggable Genome

Protein Pathways: Chemokine signaling pathway, Focal adhesion, Leukocyte transendothelial migration, Long-

term potentiation, MAPK signaling pathway, Neurotrophin signaling pathway, Renal cell

carcinoma

Product images:



Coomassie blue staining of purified RAP1A protein (Cat# [TP315248]). The protein was produced from HEK293T cells transfected with RAP1A cDNA clone (Cat# [RC215248]) using MegaTran 2.0 (Cat# [TT210002]).