

Product datasheet for TP315267L

HDAC9 (NM_014707) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human histone deacetylase 9 (HDAC9), transcript variant 3, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC215267 representing NM_014707 Red =Cloning site Green =Tags(s)

MHSMISSVDVKSEVPVGLPEISPLDLRTDLRMMMPVDPVREKQLQQEELLIQQQQIQKQLLIAEFQK
QHENLTRQHQAQLQEHIKELLAIKQQELLEKEQKLEQQRQEVEVERHRREQQLPPLRGKDRGRERAVAS
TEVKQKLQEFLLSKSATKDTPTNGKNHSVSRHPKLWYTAHHSTLDQSSPPLSGTSPSYKYTLPGAQDAK
DDFPLRKTAEPNLKVRSLKQKVAERRSSPLLRRKDGNNVTSFKKRMFEVTESSVSSSPGSGPSSPNN
GPTGSVTENETSVLPPTPHAEQMVSQQRILIHEDSMNLLSLYTSPLPNITLGLPAVPSQLNASNSLKEK
QKCETQTLRQGVPLPGQYGGSSIPASSSHPHVTLEGKPPNSSHQALLQHLLLKEQMRQQKLLVAGGVPLHP
QSPLATKERISPGIRGTHKLPRHRPLNRTQSAPLPQSTLAQLVIQQQHQQFLEKQKQYQQQIHMNKLLSK
SIEQLKQPGSHLEEAEEELQGDQAMQEDRAPSSGNSTRSDSSACVDDTLGQVGVAVKVKKEEPVDSDEDAQI
QEMESGEQA AFM QQVIGKDLAPGFVIKVII

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	65.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_055522
Locus ID:	9734
UniProt ID:	Q9UKV0
RefSeq Size:	4238
Cytogenetics:	7p21.1
RefSeq ORF:	1770
Synonyms:	HD7; HD7b; HD9; HDAC; HDAC7; HDAC7B; HDAC9B; HDAC9FL; HDRP; MITR
Summary:	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene has sequence homology to members of the histone deacetylase family. This gene is orthologous to the <i>Xenopus</i> and mouse MITR genes. The MITR protein lacks the histone deacetylase catalytic domain. It represses MEF2 activity through recruitment of multicomponent corepressor complexes that include CtBP and HDACs. This encoded protein may play a role in hematopoiesis. Multiple alternatively spliced transcripts have been described for this gene but the full-length nature of some of them has not been determined. [provided by RefSeq, Jul 2008]
Protein Families:	Druggable Genome, Transcription Factors

Product images:



Coomassie blue staining of purified HDAC9 protein (Cat# [TP315267]). The protein was produced from HEK293T cells transfected with HDAC9 cDNA clone (Cat# [RC215267]) using MegaTran 2.0 (Cat# [TT210002]).