

Product datasheet for TP315363M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Livin (BIRC7) (NM_022161) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human baculoviral IAP repeat-containing 7 (BIRC7), transcript variant 2,

100 µg

Species: Human Expression Host: HEK293T

Expression cDNA >RC215363 representing NM_022161 Clone or AA Sequence: Red=Cloning site Green=Tags(s)

MGPKDSAKCLHRGPQPSHWAAGDGPTQERCGPRSLGSPVLGLDTCRAWDHVDGQILGQLRPLTEEEEEG AGATLSRGPAFPGMGSEELRLASFYDWPLTAEVPPELLAAAGFFHTGHQDKVRCFFCYGGLQSWKRGDDP WTEHAKWFPSCQFLLRSKGRDFVHSVQETHSQLLGSWDPWEEPEDAAPVAPSVPASGYPELPTPRREVQS ESAQEPGARDVEAQLRRLQEERTCKVCLDRAVSIVFVPCGHLVCAECAPGLQLCPICRAPVRSRVRTFLS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 30.7 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling

conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 071444

Locus ID: 79444



Livin (BIRC7) (NM_022161) Human Recombinant Protein - TP315363M

UniProt ID: Q96CA5

RefSeq Size: 1268

Cytogenetics: 20q13.33

RefSeq ORF: 840

Synonyms: KIAP; LIVIN; ML-IAP; MLIAP; RNF50


Summary: This gene encodes a member of the inhibitor of apoptosis protein (IAP) family, and contains a

single copy of a baculovirus IAP repeat (BIR) as well as a RING-type zinc finger domain. The BIR domain is essential for inhibitory activity and interacts with caspases, while the RING finger domain sometimes enhances antiapoptotic activity but does not inhibit apoptosis alone. Elevated levels of the encoded protein may be associated with cancer progression and play a role in chemotherapy sensitivity. Alternative splicing results in multiple transcript variants

[provided by RefSeq, Jul 2013]

Protein Families: Druggable Genome

Product images:



Coomassie blue staining of purified BIRC7 protein (Cat# [TP315363]). The protein was produced from HEK293T cells transfected with BIRC7 cDNA clone (Cat# [RC215363]) using MegaTran 2.0 (Cat# [TT210002]).