

Product datasheet for **TP315432M**

TCF3 / E2A (TCF3) (NM_003200) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human transcription factor 3 (E2A immunoglobulin enhancer binding factors E12/E47) (TCF3), transcript variant 1, 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC215432 representing NM_003200 Red =Cloning site Green =Tags(s)

MNQPQRMAPVGTDKELSDLLDFSMMFPLPVTNGKGRPASLAGAQFGGSGLEDRPSSGSWGSGDQSSSS
FD
PSRTFSEGTHFTESHSSLSSTFLGPGLGGKSGERGAYASFGRDAGVGGLTQAGFLSGELALNSPGPLSP
SGMKGTSQYYPSYSGSSRRRAADGSLDTQPKKVRKVPPGLPSSVYPPSSGEDYGRDATAYPSAKTPSSTY
PAPFYVADGSLHPSAELWSPPGQAGFGPMLGGGSSPLPLPPGSGPVGSSGSSSTFGGLHQHERMGYQL
HG
AEVNGGLPSASSFSSAPGATYGGVSSHTPPVSGADSLGSRGTTAGSSGDALGKALASIYSPDHSSNNFS
SSPSTPVGSPQGLAGTSQWPRAGAPGALSPSYDGGLHGLQSKIEDHLDEAIHVLRSHAVGTAGDMHTLLP
GHGALASGFTSPMSLGGRHAGLVGGSHPEDGLAGSTSLMHNHAALPSQPGTLPDLRPPDSYSGLGRAG
A
TAAASEIKREEKEDEENTSAADHSEEEKKELKAPRARTSPDEDEDLLPPEQKAEREKERRVANNARERL
RVRDINEAFKELGRMCQLHLNSEKPQTKLLILHQAVSVILNLEQQVRERNLNPKAACLKRREEEKVSGVW
GDPQMVLSAPHPGLSEAHNPAGHM


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	67.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.



[View online »](#)

Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_003191
Locus ID:	6929
UniProt ID:	P15923
RefSeq Size:	4396
Cytogenetics:	19p13.3
RefSeq ORF:	1962
Synonyms:	AGM8; bHLHb21; E2A; E47; ITF1; p75; TCF-3; VDIR
Summary:	This gene encodes a member of the E protein (class I) family of helix-loop-helix transcription factors. E proteins activate transcription by binding to regulatory E-box sequences on target genes as heterodimers or homodimers, and are inhibited by heterodimerization with inhibitor of DNA-binding (class IV) helix-loop-helix proteins. E proteins play a critical role in lymphopoiesis, and the encoded protein is required for B and T lymphocyte development. Deletion of this gene or diminished activity of the encoded protein may play a role in lymphoid malignancies. This gene is also involved in several chromosomal translocations that are associated with lymphoid malignancies including pre-B-cell acute lymphoblastic leukemia (t(1;19), with PBX1), childhood leukemia (t(19;19), with TFPT) and acute leukemia (t(12;19), with ZNF384). Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and a pseudogene of this gene is located on the short arm of chromosome 9. [provided by RefSeq, Sep 2011]
Protein Families:	Druggable Genome, ES Cell Differentiation/IPS, Induced pluripotent stem cells, Transcription Factors

Product images:

Coomassie blue staining of purified TCF3 protein (Cat# [TP315432]). The protein was produced from HEK293T cells transfected with TCF3 cDNA clone (Cat# [RC215432]) using MegaTran 2.0 (Cat# [TT210002]).