

Product datasheet for TP315520

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Morg1 (WDR83) (NM 001099737) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Homo sapiens mitogen-activated protein kinase organizer 1

(MORG1), transcript variant 1, 20 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC215520 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MAFPEPKPRPPELPQKRLKTLDCGQGAVRAVRFNVDGNYCLTCGSDKTLKLWNPLRGTLLRTYSGHGYEV LDAAGSFDNSSLCSGGGDKAVVLWDVASGQVVRKFRGHAGKVNTVQFNEEATVILSGSIDSSIRCWDCRS RRPEPVQTLDEARDGVSSVKVSDHEILAGSVDGRVRRYDLRMGQLFSDYVGSPITCTCFSRDGQCTLVSS LDSTLRLLDKDTGELLGEYKGHKNQEYKLDCCLSERDTHVVSCSEDGKVFFWDLVEGALALALPVGSGVV

QSLAYHPTEPCLLTAMGGSVQCWREEAYEAEDGAG

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 34.2 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeg: NP 001093207

Locus ID: 84292





UniProt ID: Q9BRX9

RefSeq Size: 1476

Cytogenetics: 19p13.13

RefSeq ORF: 945


Synonyms: MORG1

Summary: This gene encodes a member of the WD-40 protein family. The protein is proposed to

function as a molecular scaffold for various multimeric protein complexes. The protein associates with several components of the extracellular signal-regulated kinase (ERK) pathway, and promotes ERK activity in response to serum or other signals. The protein also interacts with egl nine homolog 3 (EGLN3, also known as PHD3) and regulates expression of hypoxia-inducible factor 1, and has been purified as part of the spliceosome. Alternative

splicing results in multiple transcript variants. [provided by RefSeq, Oct 2009]

Product images:



Coomassie blue staining of purified WDR83 protein (Cat# TP315520). The protein was produced from HEK293T cells transfected with WDR83 cDNA clone (Cat# [RC215520]) using MegaTran 2.0 (Cat# [TT210002]).