

## Product datasheet for **TP315618L**

### ATP6V0E1 (NM\_003945) Human Recombinant Protein

#### Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human ATPase, H <sup>+</sup> transporting, lysosomal 9kDa, V0 subunit e1 (ATP6V0E1), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC215618 representing NM_003945 <b>Red</b> =Cloning site <b>Green</b> =Tags(s)  MAYHGLTVPLIVMSVFWGFGVGLVPWFIPKGPNRGVIITMLVTCSVCCYLFWLIAILAQLNPLFGPQLKN ETIWYLKYHWP  <b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b>
Tag:	C-Myc/DDK
Predicted MW:	9.2 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<a href="#">NP_003936</a>
Locus ID:	8992
UniProt ID:	<a href="#">O15342</a>
RefSeq Size:	894



[View online »](#)

<b>Cytogenetics:</b>	5q35.1
<b>RefSeq ORF:</b>	243
<b>Synonyms:</b>	ATP6H; ATP6V0E; M9.2; Vma21; Vma21p
<b>Summary:</b>	<p>This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'', and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This encoded protein is possibly part of the V0 subunit. Since two nontranscribed pseudogenes have been found in dog, it is possible that the localization to chromosome 2 for this gene by radiation hybrid mapping is representing a pseudogene. Genomic mapping puts the chromosomal location on 5q35.3. [provided by RefSeq, Jul 2008]</p>
<b>Protein Families:</b>	Transmembrane
<b>Protein Pathways:</b>	Epithelial cell signaling in Helicobacter pylori infection, Metabolic pathways, Oxidative phosphorylation, Vibrio cholerae infection

### Product images:



Coomassie blue staining of purified ATP6V0E1 protein (Cat# [TP315618]). The protein was produced from HEK293T cells transfected with ATP6V0E1 cDNA clone (Cat# [RC215618]) using MegaTran 2.0 (Cat# [TT210002]).