

Product datasheet for **TP315903**

SARM (SARM1) (NM_015077) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human sterile alpha and TIR motif containing 1 (SARM1), 20 µg

Species: Human

Expression Host: HEK293T

Expression cDNA Clone >RC215903 representing NM_015077

or AA Sequence: **Red**=Cloning site **Green**=Tags(s)

MVLTLILLSAYKLCRFFAMSGPRPGAERLAVPGPDGGGGTGPWWAAGGRGPREVSPGAGTEVQDALERALP
ELQQALSALKQAGGARAVGAGLAEVFQLVEEAWLLPAVGREVAQGLCDAIRLDGGDLLLRLLQAPELET
RVQAARLLEQILVAENRDRVARIGLVILNLAKEREPVELARSVAGILEHMFKHSEETCQRLVAAGGLDA
VLYWCRRTDPALLRHCALALGNALHGGQAVQRRMVEKRAAEWLFPLAFSKEDELLRLHACLAVAVLATN
KEVEREVERSGTLALVEPLVASLDPGRFARCLVDASDTSQGRGPDDLQRLVPLLDNSRLEAQCIGAFYLC
AEAAIKSLQGKTKVFSDIGAIQSLKRLVSYSTNGTKSALAKRALRLLGEEVPRPILPSVPSWKEAEVQTW
LQQIGFSKYCESFREQQVDGDLRLTEELQTDLGMKSGITRKRFFRELTELKTFANYSTCDRSNLADW
LGS�DPRFRQYTYGLVSCGLDRSLLHRVSEQQLEDCGIHLGVHRARILTAAREMLHSPLPCTGGKPSGD
TPDVFISYRRNSGSQLASLLKVHLQLHGFSVFIDVEKLEAGKFEDKLIQSVMGARNFVLVSPGALDKCM
QDHDCKDWVHKEIVTALSCGKNIVPIIDGFEWPEPQVLPEDMQAVLTFNGIKWSHEYQEATIEKIIRFLQ
GRSSRDSSAGSDTSLEGAAPMGPT

SGPTRRRLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 79.2 kDa

Concentration: >0.1 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.


Storage: Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_055892
Locus ID:	23098
UniProt ID:	Q6SZW1 , Q0D2N8 , Q05B42
RefSeq Size:	7001
Cytogenetics:	17q11.2
RefSeq ORF:	2172
Synonyms:	hSARM1; HsTIR; MyD88-5; SAMD2; SARM
Summary:	Negative regulator of MYD88- and TRIF-dependent toll-like receptor signaling pathway which plays a pivotal role in activating axonal degeneration following injury. Promotes Wallerian degeneration an injury-induced axonal death pathway which involves degeneration of an axon distal to the injury site. Can activate neuronal death in response to stress. Regulates dendritic arborization through the MAPK4-JNK pathway. Involved in innate immune response. Inhibits both TICAM1/TRIF- and MYD88-dependent activation of JUN/AP-1, TRIF-dependent activation of NF-kappa-B and IRF3, and the phosphorylation of MAPK14/p38.[UniProtKB/Swiss-Prot Function]
Protein Families:	Druggable Genome

Product images:



Coomassie blue staining of purified SARM1 protein (Cat# TP315903). The protein was produced from HEK293T cells transfected with SARM1 cDNA clone (Cat# [RC215903]) using MegaTran 2.0 (Cat# [TT210002]).