

Product datasheet for TP316020M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Histone H2A Bbd (H2AFB1) (NM_001017990) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human H2A histone family, member B1 (H2AFB1), 100 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC216020 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MPRRRRRGSSGAGGRGRTCSRTVRAELSFSVSQVERSLREGHYAQRLSRTAPVYLAAVIEYLTAKVLEL

AGNEAQNSGERNITPLLLDMVVHNDRLLSTLFNTTTISQVAPGED

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

Tag: C-Myc/DDK

Predicted MW: 12.5 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001017990

 Locus ID:
 474382

 UniProt ID:
 P0C5Y9

 RefSeq Size:
 517

Cytogenetics: Xq28





RefSeq ORF: 345

Synonyms: H2A.B; H2A.Bbd; H2AFB1


Summary: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the

chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a replication-independent histone that is a member of the histone H2A family. This gene is part of a region that is repeated three times on chromosome X, once in intron 22 of the F8 gene and twice closer to the Xq telomere. This record represents the most centromeric copy which is in intron 22 of the F8 gene. [provided by RefSeq, Oct

2015]

Protein Pathways: Systemic lupus erythematosus

Product images:



Coomassie blue staining of purified H2AFB1 protein (Cat# [TP316020]). The protein was produced from HEK293T cells transfected with H2AFB1 cDNA clone (Cat# [RC216020]) using MegaTran 2.0 (Cat# [TT210002]).