

## **Product datasheet for TP316061L**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

#### **GBA (NM 000157) Human Recombinant Protein**

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human glucosidase, beta; acid (includes glucosylceramidase) (GBA),

transcript variant 1, 1 mg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC216061 representing NM\_000157 **or AA Sequence:** Red=Cloning site Green=Tags(s)

MEFSSPSREECPKPLSRVSIMAGSLTGLLLLQAVSWASGARPCIPKSFGYSSVVCVCNATYCDSFDPPTF
PALGTFSRYESTRSGRRMELSMGPIQANHTGTGLLLTLQPEQKFQKVKGFGGAMTDAAALNILALSPPAQ
NLLLKSYFSEEGIGYNIIRVPMASCDFSIRTYTYADTPDDFQLHNFSLPEEDTKLKIPLIHRALQLAQRP
VSLLASPWTSPTWLKTNGAVNGKGSLKGQPGDIYHQTWARYFVKFLDAYAEHKLQFWAVTAENEPSAGLL
SGYPFQCLGFTPEHQRDFIARDLGPTLANSTHHNVRLLMLDDQRLLLPHWAKVVLTDPEAAKYVHGIAVH
WYLDFLAPAKATLGETHRLFPNTMLFASEACVGSKFWEQSVRLGSWDRGMQYSHSIITNLLYHVVGWTDW
NLALNPEGGPNWVRNFVDSPIIVDITKDTFYKQPMFYHLGHFSKFIPEGSQRVGLVASQKNDLDAVALMH
PDGSAVVVVLNRSSKDVPLTIKDPAVGFLETISPGYSIHTYLWRRQ

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK
Predicted MW: 55.5 kDa

**Concentration:**  $>0.05 \mu g/\mu L$  as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

#### GBA (NM\_000157) Human Recombinant Protein - TP316061L

**Bioactivity:** The enzymatic activity of TP316061 (GBA) was measured by its ability to hydrolyze a

fluorescent substrate 4-methylumbelliferyl- $\beta$ -D-glucopyranoside. The specific activity is > 70,000 pmol/hour/µg, as measured under the following conditions: 27 ng of GBA was incubated with 10 mM 4-methylumbelliferyl-  $\beta$ -D-glucopyranoside in the following buffer at 37°C for 40 min: 150 mM citrate-phosphate buffer, pH 5.4, 0.25% (w/w) sodium taurocholate, 0.25% (w/w) Triton X-100, and 1% bovine serum albumin. The reaction was terminated by adding 0.5 volume of 1M glycine buffer, pH 12.5. The hydrolyzed product of reaction, 4-methylumbelliferone (4-MU), was measured using a FlexStation 3 microplate reader (Ex365/Em445). Specific activity of GBA was calculated based on a standard curve of known

concentration of 4-MU.

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 000148

**Locus ID:** 2629

UniProt ID: <u>P04062</u>, <u>A0A068F658</u>

RefSeq Size: 2324 Cytogenetics: 1q22 RefSeq ORF: 1608

**Synonyms:** GBA1; GCB; GLUC

**Summary:** This gene encodes a lysosomal membrane protein that cleaves the beta-glucosidic linkage of

glycosylceramide, an intermediate in glycolipid metabolism. Mutations in this gene cause

Gaucher disease, a lysosomal storage disease characterized by an accumulation of

glucocerebrosides. A related pseudogene is approximately 12 kb downstream of this gene on chromosome 1. Alternative splicing results in multiple transcript variants. [provided by RefSeq,


Jan 2010]


**Protein Families:** Druggable Genome

Protein Pathways: Lysosome, Metabolic pathways, Other glycan degradation, Sphingolipid metabolism



# **Product images:**





Coomassie blue staining of purified GBA protein (Cat# [TP316061]). The protein was produced from HEK293T cells transfected with GBA cDNA clone (Cat# [RC216061]) using MegaTran 2.0 (Cat# [TT210002]).