

## Product datasheet for TP316317M

### CCBL1 (KYAT1) (NM\_004059) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human cysteine conjugate-beta lyase, cytoplasmic (CCBL1), transcript variant 1, 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC216317 representing NM_004059 Red=Cloning site Green=Tags(s)

MAKQLQARRLDGIDYNPWVEFVKLASEHDVWNLGQGFPDFPPDFAVEAFQHAVSGDFMLNQYTKTFGYP  
PLTKILASFFGELLGQEIDPLRNVLVTGGYGALFTAFQALVDEGDEVIIIPEFFDCYEPMTMMAGGRP  
VSLKPGPIQNGELGSSSNWQLDPMELAGKFTSRTKALVLNTPNPLGKVFSSREELELVASLCQQHDVVC  
ITDEVYQWMVYDGHQHISIASLPGMWERTLTIGSAGKTFATGWKVGWVLPDHPHIMKHLRTVHQNSV  
FHCPTQSQAAVAESFEREQLLFRQPSSYFVQFPQAMQRCRDHMIIRSLQSVGLKPIIPQGSYFLITDISDFK  
RKMPLDLPGAVDEPYDRRFVKWMIKNGKLVVAIPVSIFYSVPHQKHFDHYIRFCFVKDEATLQAMDEKLRK  
WKVEL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	47.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Bioactivity:	The specific activity of KATI was determined by measuring the product Kynurenic acid formation from a conversion of Kynurenine. The reaction was carried out at 37° for 15min in the buffer containing 100 mM PBS, pH7.4, 2 mM a-oxoglutarate, 40µM PLP (pyridoxal 5'-phosphate), and 0.5 mM kynurenine as the substrate.
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.




[View online »](#)

<b>Storage:</b>	Store at -80°C.
<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<u>NP_004050</u>
<b>Locus ID:</b>	883
<b>UniProt ID:</b>	<u>Q16773</u> , <u>A8K563</u>
<b>RefSeq Size:</b>	1925
<b>Cytogenetics:</b>	9q34.11
<b>RefSeq ORF:</b>	1266
<b>Synonyms:</b>	CCBL1; GTK; KAT1; KATI
<b>Summary:</b>	This gene encodes a cytosolic enzyme that is responsible for the metabolism of cysteine conjugates of certain halogenated alkenes and alkanes. This metabolism can form reactive metabolites leading to nephrotoxicity and neurotoxicity. Increased levels of this enzyme have been linked to schizophrenia. Multiple transcript variants that encode different isoforms have been identified for this gene. [provided by RefSeq, Jul 2008]

### Product images:





Coomassie blue staining of purified KYAT1 protein (Cat# [TP316317]). The protein was produced from HEK293T cells transfected with KYAT1 cDNA clone (Cat# [RC216317]) using MegaTran 2.0 (Cat# [TT210002]).