

## Product datasheet for **TP316714**

### **MAK3 (NAA30) (NM\_001011713) Human Recombinant Protein**

#### **Product data:**


<b>Product Type:</b>	Recombinant Proteins
<b>Description:</b>	Recombinant protein of human N-acetyltransferase 12 (GCN5-related, putative) (NAT12), 20 µg
<b>Species:</b>	Human
<b>Expression Host:</b>	HEK293T
<b>Expression cDNA Clone or AA Sequence:</b>	>RC216714 representing NM_001011713 <b>Red</b> =Cloning site <b>Green</b> =Tags(s)  MAEVPPGPSSLLPPPAPPAPAAVEPRCPFPAGAALACCSEDEEDDEEHEGGGSRSPAGGESATVAAKGHP CLRCPQPPQEQQQLNGLISPELRHLRAAASLKSKVLSVAEVAATTATPDGGPRATATKGAGVHSGERPPH SLSSNARTAVPSPVEAAAAADPAAARNGLAEGTEQEEEEDEQVRLSSSLTADCSLRSPSGREVEPGED RTIRYVRYESELQMPDIMRLITKDLSEPSIYTYRYFIHNWPQLCFLAMVGEECVGAIVCKLDMHKKMFR RGYIAMLAVDSKYRRNGIGTNLVKKAIYAMVEGDCDEVVLEITEITNKSALKLYENLGFVRDKRLFRYYLN GVDALRLKLWLR  <b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b>
<b>Tag:</b>	C-Myc/DDK
<b>Predicted MW:</b>	39.1 kDa
<b>Concentration:</b>	>0.05 µg/µL as determined by microplate BCA method
<b>Purity:</b>	> 80% as determined by SDS-PAGE and Coomassie blue staining
<b>Buffer:</b>	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
<b>Preparation:</b>	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
<b>Note:</b>	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
<b>Storage:</b>	Store at -80°C.
<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<u>NP_001011713</u>



[View online »](#)

Locus ID:	122830
UniProt ID:	<a href="#">Q147X3</a>
RefSeq Size:	2891
Cytogenetics:	14q22.3
RefSeq ORF:	1086
Synonyms:	C14orf35; MAK3; Mak3p; NAT12; NAT12P
Summary:	Catalytic subunit of the N-terminal acetyltransferase C (NatC) complex. Catalyzes acetylation of the N-terminal methionine residues of peptides beginning with Met-Leu-Ala and Met-Leu-Gly. Necessary for the lysosomal localization and function of ARL8B suggesting that ARL8B is a NatC substrate.[UniProtKB/Swiss-Prot Function]

### Product images:



Coomassie blue staining of purified NAA30 protein (Cat# TP316714). The protein was produced from HEK293T cells transfected with NAA30 cDNA clone (Cat# [RC216714]) using MegaTran 2.0 (Cat# [TT210002]).