

Product datasheet for TP316714L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MAK3 (NAA30) (NM_001011713) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human N-acetyltransferase 12 (GCN5-related, putative) (NAT12), 1

mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC216714 representing NM_001011713

or AA Sequence: Red=Cloning site Green=Tags(s)

MAEVPPGPSSLLPPPAPPAPAAVEPRCPFPAGAALACCSEDEEDDEEHEGGGSRSPAGGESATVAAKGHP CLRCPQPPQEQQLNGLISPELRHLRAAASLKSKVLSVAEVAATTATPDGGPRATATKGAGVHSGERPPH SLSSNARTAVPSPVEAAAASDPAAARNGLAEGTEQEEEEEDEQVRLLSSSLTADCSLRSPSGREVEPGED RTIRYVRYESELQMPDIMRLITKDLSEPYSIYTYRYFIHNWPQLCFLAMVGEECVGAIVCKLDMHKKMFR RGYIAMLAVDSKYRRNGIGTNLVKKAIYAMVEGDCDEVVLETEITNKSALKLYENLGFVRDKRLFRYYLN

GVDALRLKLWLR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 39.1 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001011713





MAK3 (NAA30) (NM_001011713) Human Recombinant Protein - TP316714L

Locus ID: 122830

 UniProt ID:
 Q147X3

 RefSeq Size:
 2891

Cytogenetics: 14q22.3 RefSeq ORF: 1086


Synonyms: C14orf35; MAK3; Mak3p; NAT12; NAT12P

Summary: Catalytic subunit of the N-terminal acetyltransferase C (NatC) complex. Catalyzes acetylation

of the N-terminal methionine residues of peptides beginning with Met-Leu-Ala and Met-Leu-Gly. Necessary for the lysosomal localization and function of ARL8B suggesting that ARL8B is

a NatC substrate.[UniProtKB/Swiss-Prot Function]

Product images:



Coomassie blue staining of purified NAA30 protein (Cat# [TP316714]). The protein was produced from HEK293T cells transfected with NAA30 cDNA clone (Cat# [RC216714]) using MegaTran 2.0 (Cat# [TT210002]).