

Product datasheet for TP316785

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL10 (NM_000572) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Homo sapiens interleukin 10 (IL10), 20 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC216785 representing NM_000572 or AA Sequence: Red=Cloning site Green=Tags(s)

MHSSALLCCLVLLTGVRASPGQGTQSENSCTHFPGNLPNMLRDLRDAFSRVKTFFQMKDQLDNLLLKES

L

LEDFKGYLGCQALSEMIQFYLEEVMPQAENQDPDIKAHVNSLGENLKTLRLRRCHRFLPCENKSKAVE

QVKNAFNKLQEKGIYKAMSEFDIFINYIEAYMTMKIRN

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 18.6 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 000563

Locus ID: 3586

UniProt ID: P22301





RefSeq Size: 1629

Cytogenetics: 1q32.1 RefSeq ORF: 534

Synonyms: CSIF; GVHDS; IL-10; IL10A; TGIF

Summary: The protein encoded by this gene is a cytokine produced primarily by monocytes and to a

lesser extent by lymphocytes. This cytokine has pleiotropic effects in immunoregulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II Ags, and costimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. This cytokine can block NF-kappa B activity, and is involved in the regulation of the JAK-STAT signaling pathway. Knockout studies in mice suggested the function of this cytokine as an essential immunoregulator in the intestinal tract. Mutations in this gene are associated with an increased susceptibility to HIV-1 infection and rheumatoid

arthritis. [provided by RefSeq, May 2020]


Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein

Protein Pathways: Allograft rejection, Asthma, Autoimmune thyroid disease, Cytokine-cytokine receptor

interaction, Jak-STAT signaling pathway, Systemic lupus erythematosus, T cell receptor

signaling pathway

Product images:



Coomassie blue staining of purified IL10 protein (Cat# TP316785). The protein was produced from HEK293T cells transfected with IL10 cDNA clone (Cat# [RC216785]) using MegaTran 2.0 (Cat# [TT210002]).