

Product datasheet for TP317022L

HCK (NM_002110) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human hemopoietic cell kinase (HCK), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC217022 representing NM_002110 Red=Cloning site Green=Tags(s)

MGGRSSCEDPGCPRDEERAPRMGCMKSKFLQVGGNTFSKTETSASPHCPVYVPDPTSTIKPGPNSHNSNT
 PGIREAGSEDIIVALDYEDIAIHEDLSFQKGDQMVVLEESGEWWKARSLATRKEGYIPSNYVARVDSLE
 TEEWFFKGISRKDAERQLLAPGNMLGSFMIRDSETTKGSYSLSVRDYDPRQGDTVKHYKIRTLDNNGFYI
 SPRSTFSTLQELVDHYKKGNDGLCQKLSVPCMSKPKQPWEKDAWEIPRESLKLEKKGAGQFGEVWMAT
 YNKHTKVAVKTMKPGSMSVEAFLAEANVMKTLQHDKLVKLHAVTKIPIYIITEFMAKGSLLDFLKSDEG
 SKQPLPKLIDFSAQIAEGMAFIEQRNYIHRDLRAANILVSASLVCKIADFGLARVIEDNEYTAREGAKFP
 IKWTAPEAINFGSFTIKSDVWSFGILLMEIVTYGRIPYPGMSNPEVIRALERGYRMPRENCPEELYNIM
 MRCWKNRPEERPTFEYIQSVLDDFYTATESQYQQQP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	59.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq: [NP_002101](#)

Locus ID: 3055

UniProt ID: [P08631](#)

RefSeq Size: 2168

Cytogenetics: 20q11.21

RefSeq ORF: 1578


Synonyms: JTK9; p59Hck; p61Hck

Summary: The protein encoded by this gene is a member of the Src family of tyrosine kinases. This protein is primarily hemopoietic, particularly in cells of the myeloid and B-lymphoid lineages. It may help couple the Fc receptor to the activation of the respiratory burst. In addition, it may play a role in neutrophil migration and in the degranulation of neutrophils. Multiple isoforms with different subcellular distributions are produced due to both alternative splicing and the use of alternative translation initiation codons, including a non-AUG (CUG) codon. [provided by RefSeq, Feb 2010]

Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: Chemokine signaling pathway, Fc gamma R-mediated phagocytosis

Product images:



Coomassie blue staining of purified HCK protein (Cat# [TP317022]). The protein was produced from HEK293T cells transfected with HCK cDNA clone (Cat# [RC217022]) using MegaTran 2.0 (Cat# [TT210002]).