

## Product datasheet for TP317103

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Glycophorin C (GYPC) (NM\_016815) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human glycophorin C (Gerbich blood group) (GYPC), transcript variant

2, 20 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC217103 representing NM\_016815

or AA Sequence: Red=Cloning site Green=Tags(s)

MWSTRSPNSTAWPLSLEPDPGMSGWPDGRMETSTPTIMDIVVIAGVIAAVAIVLVSLLFVMLRYMYRHKG

TYHTNEAKGTEFAESADAALQGDPALQDAGDSSRKEYFI

**TRTRPL**EQKLISEEDLAANDILDYKDDDDK**V** 

Tag: C-Myc/DDK

**Predicted MW:** 11.7 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 058131

 Locus ID:
 2995

 UniProt ID:
 P04921

 RefSeq Size:
 1019



Cytogenetics: 2q14.3

RefSeq ORF: 327

Synonyms: CD236; CD236R; GE; GE:GPC:GPD:GYPD; GPC; GPD; GYPD; PAS-2; PAS-2'


**Summary:** Glycophorin C (GYPC) is an integral membrane glycoprotein. It is a minor species carried by

human erythrocytes, but plays an important role in regulating the mechanical stability of red cells. A number of glycophorin C mutations have been described. The Gerbich and Yus phenotypes are due to deletion of exon 3 and 2, respectively. The Webb and Duch antigens, also known as glycophorin D, result from single point mutations of the glycophorin C gene. The glycophorin C protein has very little homology with glycophorins A and B. Alternate

splicing results in multiple transcript variants. [provided by RefSeq, Feb 2012]

**Protein Families:** Druggable Genome, Transmembrane

## **Product images:**



Coomassie blue staining of purified GYPC protein (Cat# TP317103). The protein was produced from HEK293T cells transfected with GYPC cDNA clone (Cat# [RC217103]) using MegaTran 2.0 (Cat# [TT210002]).