

## Product datasheet for **TP317536M**

### TRIM68 (NM\_018073) Human Recombinant Protein

#### Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human tripartite motif-containing 68 (TRIM68), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC217536 representing NM_018073 <b>Red</b> =Cloning site <b>Green</b> =Tags(s)
	<p>MDPTALVEAIVEEVACPICMTFLREPMSIDCGHSFCHSCLSGLWEIPGESQNWGYTCPLCRAPVQPRNLR PNWQLANVVEKVRLLRLHPGMGLKGDLCERHGEKLMFCKEDVLIMCEACSQSPEHEAHSVPMEDVA WE YKWELEHALEHLKKEQEEAWKLEVGERKRTATWKIQVETRQKSIWVEFEKYQRLLEKKQPPHRQLGAEVA AALASLQREAAETMQKLELNHSELIQQSQVLWRMIAELKERSQRPVRWMLQDIQEVNRSKSWSLQQPE P ISLELKTDCRVLGLREILKTYAADVRLDPDTAYSRLIVSEDRKRVHYGDTNQLPDNPERFYRYNIVLGS QCISSGRHYWEVEVGDRSEWGLGVCKQNVDRKEVVYLSPHYGFWVIRLRKGNEYRAGTDEYPILSLPVPP RRVGIFVDYEAHDISFYNVTDGSHIFTPRPYPFGRLLPYFSPCYSIGTNNTAPLAICSLDGED</p> <p><b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b></p>
Tag:	C-Myc/DDK
Predicted MW:	56.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<a href="#">NP_060543</a>
<b>Locus ID:</b>	55128
<b>UniProt ID:</b>	<a href="#">Q6AZZ1</a>
<b>RefSeq Size:</b>	3321
<b>Cytogenetics:</b>	11p15.4
<b>RefSeq ORF:</b>	1455
<b>Synonyms:</b>	GC109; RNF137; SS-56; SS56
<b>Summary:</b>	This gene encodes a member of the tripartite motif-containing protein family, whose members are characterized by a "really interesting new gene" (RING) finger domain, a zinc-binding B-box motif, and a coiled-coil region. Members of this family function as E3 ubiquitin ligases and are involved in a broad range of biological processes. This gene regulates the activation of nuclear receptors, such as androgen receptor, and has been implicated in development of prostate cancer cells, where its expression increases in response to a downregulation of microRNAs. In addition, this gene participates in viral defense regulation as a negative regulator of interferon-beta. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2015]
<b>Protein Families:</b>	Druggable Genome

## Product images:



Coomassie blue staining of purified TRIM68 protein (Cat# [TP317536]). The protein was produced from HEK293T cells transfected with TRIM68 cDNA clone (Cat# [RC217536]) using MegaTran 2.0 (Cat# [TT210002]).