

Product datasheet for **TP317695M**

USP2 (NM_171997) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human ubiquitin specific peptidase 2 (USP2), transcript variant 2, 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC217695 representing NM_171997 Red =Cloning site Green =Tags(s)

MRTSYTVTLPEDPPAAPFPALAKELRPRSPLSPSLLLSTFVGLLLNKAKNSKSAQGLAGLRNLGNTCFMN
SILQCLSNTRERLDYCLQRLYMRDLHHGSNAHTALVEEFAKLIQTIWTSSPNDVWSPSEFKTQIQRYAPR
FVGYNQDAQEFLRFLLDGLHNEVNRVTLRPKSNPENLDHLPDDEKGRQMWRKYLERSRIGDLFVGQL
KSSLTCTDCGYCSTVFDPFDLSLPIAKRGYPEVTLMDCMRLFTKEDVLDGDEKPTCCRCRGRKRCIKKF
SIQRFPKILVHLKRFSESRIKTSKLTTFVNFPLRDLDLREFASENTNHAVYNLYAVSNHSGTMMGGHYT
AYCRSPGTGEWHTFNDSSVTPMSSSQVRTSDAYLLFYELASPPSRM

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	45.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_741994




[View online »](#)

Locus ID: 9099
UniProt ID: [O75604](#), [O75604-4](#)
RefSeq Size: 1378
Cytogenetics: 11q23.3
RefSeq ORF: 1188
Synonyms: UBP41; USP9

Summary: This gene encodes a member of the family of de-ubiquitinating enzymes, which belongs to the peptidase C19 superfamily. The encoded protein is a ubiquitin-specific protease which is required for TNF-alpha (tumor necrosis factor alpha) -induced NF-kB (nuclear factor kB) signaling. This protein deubiquitinates polyubiquitinated target proteins such as fatty acid synthase, murine double minute 2 (MDM2), MDM4/MDMX and cyclin D1. MDM2 and MDM4 are negative regulators of the p53 tumor suppressor and cyclin D1 is required for cell cycle G1/S transition. Multiple alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Aug 2011]

Protein Families: Protease

Product images:



Coomassie blue staining of purified USP2 protein (Cat# [TP317695]). The protein was produced from HEK293T cells transfected with USP2 cDNA clone (Cat# [RC217695]) using MegaTran 2.0 (Cat# [TT210002]).