

Product datasheet for **TP317929L**

PLAUR (NM_001005376) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Homo sapiens plasminogen activator, urokinase receptor (PLAUR), transcript variant 2, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC217929 representing NM_001005376 Red =Cloning site Green =Tags(s)
	<p>MGHPPLLPLLLLHTCVPASWGLRCMQCKTNGDCRVEECALGQDLCRTTIVRLWEEGEELELVEKSCTHS EKTNRTLSYRTGLKITSLEVVCGLDLCNQGNSGRAVTYSRSRYLEICISGSSDMSCERGRHQSLQCRSP EEQCLDVVTHWIQEGEEGRPKDDRHLRGCGYLPGCPGNGFHNNDTFHFLKCCNTTKCNEGPILELENLP QNGRQCYSCKGNSTHGCSSEETFLIDCRGPMNQCLVATGOTHERSLWGSWLPCKSTTALRPPCCEEAQATH V</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-Myc/DDK
Predicted MW:	28.9 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_001005376
Locus ID:	5329



[View online »](#)

UniProt ID: [Q03405](#)

RefSeq Size: 1437

Cytogenetics: 19q13.31

RefSeq ORF: 843


Synonyms: CD87; U-PAR; UPAR; URKR

Summary: This gene encodes the receptor for urokinase plasminogen activator and, given its role in localizing and promoting plasmin formation, likely influences many normal and pathological processes related to cell-surface plasminogen activation and localized degradation of the extracellular matrix. It binds both the proprotein and mature forms of urokinase plasminogen activator and permits the activation of the receptor-bound pro-enzyme by plasmin. The protein lacks transmembrane or cytoplasmic domains and may be anchored to the plasma membrane by a glycosyl-phosphatidylinositol (GPI) moiety following cleavage of the nascent polypeptide near its carboxy-terminus. However, a soluble protein is also produced in some cell types. Alternative splicing results in multiple transcript variants encoding different isoforms. The proprotein experiences several post-translational cleavage reactions that have not yet been fully defined. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Complement and coagulation cascades

Product images:



Coomassie blue staining of purified PLAUR protein (Cat# [TP317929]). The protein was produced from HEK293T cells transfected with PLAUR cDNA clone (Cat# [RC217929]) using MegaTran 2.0 (Cat# [TT210002]).