

Product datasheet for **TP318441M**

NSMase2 (SMPD3) (NM_018667) Human Recombinant Protein


Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Homo sapiens sphingomyelin phosphodiesterase 3, neutral membrane (neutral sphingomyelinase II) (SMPD3), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC218441 representing NM_018667 Red =Cloning site Green =Tags(s)
	MVLYTTPFPNSCLSALHCVSWALIFPCYWLVDRLAASFIPTTYEKQRADDPCCQLLCTALFTPIYLAL LVASLPFAFLGFLFWSPQLSARRPYIYSRLEDKGLAGGAALLSEWKGTPGKSFCEFATANVCLLPDSLAR VNNLFNTQARAKEIGQRIRNGAARPQIKIYIDSPNTSISAASFSSLVSPQGGDGVARAVPGSIKRTASV EYKGDGGRHPGDEAANGPASGDPVDSSSPEDACIVRIGGEEGGRPPEADDPVPGGQARNGAGGGPRGQTP NHNQQDGDGSGSLGSPSASRESLVKGRAGPDTASGEPGANSKLLYKASVKKAAARRRRHPDEAFDHEVS AFFPANLDFLCLQEVEFDKRAATKLKEQLHGFEYILYDVG VYGCQGCCSFKCLNSGLLFASRYPIMDVAY HCYPNKCND DALASKGALFLKVQVGSTPQDQRIVGYIACHTLHAPQEDSAIRCGQLDLLQDWLADFRKST SSSSAANPEELVAFDVVCGDFNFDNCSSDDKLEQQHSLFTHYRDP CRLGPGEEKPWAIGTLLDNTGLYDE DVCTPDNLQK VLESEEGRREYLAFTSKSSGQKGRKELLKGNRRIDYMLHAE EGLCPDWKAEVEEFSFI TQLSGLTDHLPVAMRLMVSSGEEEA
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	70.9 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_061137
Locus ID:	55512
UniProt ID:	Q9NY59 , A8K0T6
RefSeq Size:	5284
Cytogenetics:	16q22.1
RefSeq ORF:	1965
Synonyms:	NSMASE2
Summary:	Catalyzes the hydrolysis of sphingomyelin to form ceramide and phosphocholine. Ceramide mediates numerous cellular functions, such as apoptosis and growth arrest, and is capable of regulating these 2 cellular events independently. Also hydrolyzes sphingosylphosphocholine. Regulates the cell cycle by acting as a growth suppressor in confluent cells. Probably acts as a regulator of postnatal development and participates in bone and dentin mineralization. [UniProtKB/Swiss-Prot Function]
Protein Families:	Transmembrane
Protein Pathways:	Metabolic pathways, Sphingolipid metabolism

Product images:

Coomassie blue staining of purified SMPD3 protein (Cat# [TP318441]). The protein was produced from HEK293T cells transfected with SMPD3 cDNA clone (Cat# [RC218441]) using MegaTran 2.0 (Cat# [TT210002]).