

## Product datasheet for **TP318468M**

### DDI2 (NM\_032341) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human DDI1, DNA-damage inducible 1, homolog 2 ( <i>S. cerevisiae</i> ) (DDI2), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC218468 representing NM_032341 <b>Red</b> =Cloning site <b>Green</b> =Tags(s)  MLLTVYCVRRDLSEVTFSLQVDADFELHNFALCELESGIPAAESQIVYAERPLTDNHRSLASYGLKDGD VVILRQKENADPRPPVQFPNLPRIDFSSIAVPGTSSPRQRQPPGTQQSHSSPGEITSSPQGLDNPALLRD MLLANPHELSLLKERNPPLAEALLSGDLEKFSRVLVEQQQDRARREQERIRLFSADPFDLEAQAKIEEDI RQQNIEENMTIAMEEAPESFGQVVMPLYINCKVNGHPVKAFVDSGAQMTIMSQACAERCNIMRLVDRRW AG IAKGVGTQKIIGRVHLAQVQIEGDFLPCFSFISILEEQPMDMLLGLDMLKRHQCSIDLKKNVLVIGTTGSQT TFLPEGELPECARLAYGAGREDVRPEEADQELAEALQKSAEDAERQKP  <b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b>
Tag:	C-Myc/DDK
Predicted MW:	44.3 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.


[View online »](#)

RefSeq: [NP\\_115717](#)

Locus ID: 84301

UniProt ID: [Q5TDH0](#)

RefSeq Size: 1759


Cytogenetics: 1p36.21

RefSeq ORF: 1197

**Summary:** Aspartic protease that mediates the cleavage of NFE2L1/NRF1 at 'Leu-104', thereby promoting release of NFE2L1/NRF1 from the endoplasmic reticulum membrane (PubMed:27676298, PubMed:27528193). Ubiquitination of NFE2L1/NRF1 is a prerequisite for cleavage, suggesting that DDI2 specifically recognizes and binds ubiquitinated NFE2L1/NRF1 (PubMed:27528193). Seems to act as a proteasomal shuttle which links the proteasome and replication fork proteins like RTF2 (Probable). Required, with DDI1, for cellular survival following replication stress. Together or redundantly with DDI1, removes RTF2 from stalled forks to allow cell cycle progression after replication stress and maintains genome integrity (PubMed:29290612). [UniProtKB/Swiss-Prot Function]

**Protein Families:** Druggable Genome

## Product images:



Coomassie blue staining of purified DDI2 protein (Cat# [TP318468]). The protein was produced from HEK293T cells transfected with DDI2 cDNA clone (Cat# [RC218468]) using MegaTran 2.0 (Cat# [TT210002]).