

Product datasheet for **TP318536L**

HDAC10 (NM_032019) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human histone deacetylase 10 (HDAC10), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC218536 representing NM_032019 Red =Cloning site Green =Tags(s)

MGTALVYHEDMTATRLLWDDPECEIERPERLTAALDRLRQRGLEQRCLRLSAREASEEELGLVHSPEYVS
LVRETQVLGKEELQALSGQFDIYFHPSTFHCARLAAGAGLQLVDAVLTGAVQNGLALVRPPGHHGQRAA
ANGFCVFNNVAIAAAHAKQKHGLHRILVVDWDVHHGQGIQYLFEDDPSVLVYFSWHRYEHGRFWPFLRES
ADAVGRGQGLGFTVNLPNQVGMGNADYVAAFHLLLPLAFEFDPPELVLSAGFDSAIGDPEGQMQATPE
CFAHLTQLLQVLGGRVCAVLEGGYHLESLSAESVCMTVQTLLGDPAPPLSGPMAPCQSALESIQSARAAQ
APHWKSLLQQDVTAVPMSSSHSPGRPPPLPGGPVCKAAASAPSSLLDQPCLCPAPSVRTAVALTTPD
ITLVLPDVIQQEASALRETEAWARPHESLAREEALTAGKLLYLLDGMLDGQVNSGIAATPASAAAAT
LDVAVRRGLSHGAQRLLCVALGQLDRPPDLAHDGRSLWLNIRGKEAAALSMFHVSTPLPVMVTGGFLSCIL
GLVLPLAYGFQPDLLVALGPGHGLQGPHAALLAAMLRLAGGRVLALLEENSTPQLAGILARVLNGEAP
PSLGPSSVASPEDVQALMYLRGQLEPQWKMLQCHPHLVA

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	71.3 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_114408](#)

Locus ID: 83933

UniProt ID: [Q969S8](#)

RefSeq Size: 2702

Cytogenetics: 22q13.33


RefSeq ORF: 2007

Synonyms: HD10

Summary: The protein encoded by this gene belongs to the histone deacetylase family, members of which deacetylate lysine residues on the N-terminal part of the core histones. Histone deacetylation modulates chromatin structure, and plays an important role in transcriptional regulation, cell cycle progression, and developmental events. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug 2011]

Protein Families: Druggable Genome, Transcription Factors

Product images:



Coomassie blue staining of purified HDAC10 protein (Cat# [TP318536]). The protein was produced from HEK293T cells transfected with HDAC10 cDNA clone (Cat# [RC218536]) using MegaTran 2.0 (Cat# [TT210002]).