

## **Product datasheet for TP318707M**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## ARHGAP8 (NM\_181335) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human Rho GTPase activating protein 8 (ARHGAP8), transcript variant

2, 100 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone or AA Sequence:

>RC218707 protein sequence Red=Cloning site Green=Tags(s)

MAGQDPALSTSHPFYDVARHGILQVAGDDRFGRRVVTFSCCRMPPSHELDHQRLLEYLKYTLDQYVENDY TIVYFHYGLNSRNKPSLGWLQSAYKEFDRKYKKNLKALYVVHPTSFIKVLWNILKPLISHKFGKKVIYFN YLSELHEHLKYDQLVIPPEVLRYDEKLQSLHEGRTPPPTKTPPPRPPLPTQQFGVSLQYLKDKNQGELIP PVLRFTVTYLREKGLRTEGLFRRSASVQTVREIQRLYNQGKPVNFDDYGDIHIPAVILKTFLRELPQPLL TFQAYEQILGITCVESSLRVTRCRQILRSLPEHNYVVLRYLMGFLHAVSRESIFNKMNSSNLACVFGLNL IWPSQGVSSLSALVPLNMFTELLIEYYEKIFSTPEAPGEHGLAPWEQGSRAAPLQEAVPRTQATGLTKPT

LPPSPLMAARRRL

**TRTRPL**EQKLISEEDLAANDILDYKDDDDK**V** 

Tag: C-Myc/DDK
Predicted MW: 49.6 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.





**RefSeq:** NP 851852

**Locus ID:** 23779

UniProt ID: <u>P85298</u>, <u>Q6PJW1</u>

RefSeq Size: 1632

Cytogenetics: 22q13.31

RefSeg ORF: 1299


Synonyms: BPGAP1; PP610

Summary: This gene encodes a member of the RHOGAP family. GAP (GTPase-activating) family proteins

participate in signaling pathways that regulate cell processes involved in cytoskeletal changes. GAP proteins alternate between an active (GTP-bound) and inactive (GDP-bound) state based on the GTP:GDP ratio in the cell. This family member is a multidomain protein that functions to promote Erk activation and cell motility. Alternative splicing results in multiple transcript variants. Read-through transcripts from the upstream proline rich 5, renal (PRR5) gene into this gene also exist, which led to the original description of PRR5 and ARHGAP8 being a single

gene. [provided by RefSeq, Nov 2010]

## **Product images:**



Coomassie blue staining of purified ARHGAP8 protein (Cat# [TP318707]). The protein was produced from HEK293T cells transfected with ARHGAP8 cDNA clone (Cat# [RC218707]) using MegaTran 2.0 (Cat# [TT210002]).