

Product datasheet for TP318795M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

UBE2D2 (NM_003339) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human ubiquitin-conjugating enzyme E2D 2 (UBC4/5 homolog, yeast)

(UBE2D2), transcript variant 1, 100 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC218795 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MALKRIHKELNDLARDPPAQCSAGPVGDDMFHWQATIMGPNDSPYQGGVFFLTIHFPTDYPFKPPKVAF

Τ

TRIYHPNINSNGSICLDILRSQWSPALTISKVLLSICSLLCDPNPDDPLVPEIARIYKTDREKYNRIARE

WTQKYAM

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 16.6 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 003330

Locus ID: 7322



UBE2D2 (NM_003339) Human Recombinant Protein - TP318795M

UniProt ID: P62837
RefSeq Size: 2715
Cytogenetics: 5q31.2
RefSeq ORF: 441

Synonyms: E2(17)KB2; PUBC1; UBC4; UBC4/5; UBCH4; UBCH5B


Summary: Regulated degradation of misfolded, damaged or short-lived proteins in eukaryotes occurs

via the ubiquitin (Ub)-proteasome system (UPS). An integral part of the UPS system is the ubiquitination of target proteins and covalent linkage of Ub-containing proteins to form polymeric chains, marking them as targets for 26S proteasome-mediated degradation. Ubiquitination of proteins is mediated by a cascade of enzymes which includes E1 (ubiquitin activating), E2 (ubiquitin conjugating), and E3 (ubiquitin ligases) enzymes. This gene encodes a member of the E2 enzyme family. Substrates of this enzyme include the tumor suppressor protein p53 and peroxisomal biogenesis factor 5 (PEX5). Alternative splicing results in

multiple transcript variants of this gene. [provided by RefSeq, May 2013]

Protein Pathways: Ubiquitin mediated proteolysis

Product images:



Coomassie blue staining of purified UBE2D2 protein (Cat# [TP318795]). The protein was produced from HEK293T cells transfected with UBE2D2 cDNA clone (Cat# [RC218795]) using MegaTran 2.0 (Cat# [TT210002]).