

Product datasheet for TP318818L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

JAML (NM 001098526) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human adhesion molecule, interacts with CXADR antigen 1 (AMICA1),

transcript variant 1, 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC218818 representing NM 001098526

or AA Sequence: Red=Cloning site Green=Tags(s)

MFCPLKLILLPVLLDYSLGLNDLNVSPPELTVHVGDSALMGCVFQSTEDKCIFKIDWTLSPGEHAKDEYV LYYYSNLSVPIGRFQNRVHLMGDILCNDGSLLLQDVQEADQGTYICEIRLKGESQVFKKAVVLHVLPEEP KELMVHVGGLIQMGCVFQSTEVKHVTKVEWIFSGRRAKEEIVFRYYHKLRMSVEYSQSWGHFQNRVNLVG DIFRNDGSIMLQGVRESDGGNYTCSIHLGNLVFKKTIVLHVSPEEPRTLVTPAALRPLVLGGNQLVIIVG IVCATILLLPVLILIVKKTCGNKSSVNSTVLVKNTKKTNPEIKEKPCHFERCEGEKHIYSPIIVREVIEE

EEPSEKSEATYMTMHPVWPSLRSDRNNSLEKKSGGGMPKTQQAF

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 44.2 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001091996





Locus ID: 120425

UniProt ID: Q86YT9, B3KUI3

RefSeq Size: 1957 Cytogenetics: 11q23.3 RefSeq ORF: 1182

Synonyms: AMICA; AMICA1; CREA7-1; CREA7-4; Gm638


Summary: Transmembrane protein of the plasma membrane of leukocytes that control their migration

and activation through interaction with CXADR, a plasma membrane receptor found on adjacent epithelial and endothelial cells. The interaction between both receptors mediates the activation of gamma-delta T-cells, a subpopulation of T-cells residing in epithelia and involved in tissue homeostasis and repair. Upon epithelial CXADR-binding, JAML induces downstream cell signaling events in gamma-delta T-cells through PI3-kinase and MAP kinases. It results in proliferation and production of cytokines and growth factors by T-cells that in turn stimulate epithelial tissues repair. It also controls the transmigration of leukocytes within epithelial and endothelial tissues through adhesive interactions with epithelial and endothelial CXADR.

[UniProtKB/Swiss-Prot Function]

Protein Families: Transmembrane

Product images:



Coomassie blue staining of purified JAML protein (Cat# [TP318818]). The protein was produced from HEK293T cells transfected with JAML cDNA clone (Cat# [RC218818]) using MegaTran 2.0 (Cat# [TT210002]).