

## Product datasheet for **TP318920**

### Glucokinase (GCK) (NM\_033508) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human glucokinase (hexokinase 4) (GCK), transcript variant 3, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC218920 representing NM_033508 <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MPRPRSQLPQPNSQVEQILAEFQLQEEDLKKVMRRMQKEMDRGLRLETHEEASVKMLPTYVVRSTPEGSEV  
GDFLSLDLGGTNFRVMLVKVGEEGEQWSVTKKHQMY SIPEDAMTGTAEMLFDYISECISDFLDKHKMKH  
KKLPLGFTFSFPVRHEDIDKGILLNWTKGFKASGAEGNNVGLLRDAIKRRGDFEMDVAMVNDTVATMI  
SCYYEDHQCEVGMIVGTGCNACYMEEMQNVELVEGDEGRMCVNTEWGAFGDSGELDEFLLDYDRLVDESS  
ANPGQQLYEKLIGGKYMGEVLVRLVLLRLVDENLLFHGEASEQLRTRGFATRFVSVQVESDTGDRKQIYNI  
LSTLGLRPSTTDCDIVRRACESVSTRAAHMCSAGLAGVINRMRESRSEDVMRITVGDGSVYKLPSPFKE  
RFHASVRRLLTPSCEITFIESEEGSGRGAALVSAVACKKACMLGQ

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**

Tag:	C-Myc/DDK
Predicted MW:	51.9 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<a href="#">NP_277043</a>



[View online »](#)

Locus ID: 2645

UniProt ID: [P35557](#), [P35557-3](#)

RefSeq Size: 2566

Cytogenetics: 7p13

RefSeq ORF: 1392


Synonyms: FGQTL3; GK; GLK; HHF3; HK4; HKIV; HXKP; LGLK; MODY2; PNDM1

**Summary:** This gene encodes a member of the hexokinase family of proteins. Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose metabolism pathways. In contrast to other forms of hexokinase, this enzyme is not inhibited by its product glucose-6-phosphate but remains active while glucose is abundant. The use of multiple promoters and alternative splicing of this gene result in distinct protein isoforms that exhibit tissue-specific expression in the pancreas and liver. In the pancreas, this enzyme plays a role in glucose-stimulated insulin secretion, while in the liver, this enzyme is important in glucose uptake and conversion to glycogen. Mutations in this gene that alter enzyme activity have been associated with multiple types of diabetes and hyperinsulinemic hypoglycemia. [provided by RefSeq, Aug 2017]

**Protein Families:** Druggable Genome

**Protein Pathways:** Amino sugar and nucleotide sugar metabolism, Galactose metabolism, Glycolysis / Gluconeogenesis, Insulin signaling pathway, Maturity onset diabetes of the young, Metabolic pathways, Starch and sucrose metabolism, Type II diabetes mellitus

### Product images:



Coomassie blue staining of purified GCK protein (Cat# TP318920). The protein was produced from HEK293T cells transfected with GCK cDNA clone (Cat# [RC218920]) using MegaTran 2.0 (Cat# [TT210002]).