

## **Product datasheet for TP319137**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## UBXN2B (NM\_001077619) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human UBX domain protein 2B (UBXN2B), 20 μg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC219137 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MAEGGGPEPGEQERRSSGPRPPSARDLQLALAELYEDEVKCKSSKSNRPKATVFKSPRTPPQRFYSSEHE YSGLNIVRPSTGKIVNELFKEAREHGAVPLNEATRASGDDKSKSFTGGGYRLGSSFCKRSEYIYGENQLQ DVQILLKLWSNGFSLDDGELRPYNEPTNAQFLESVKRGEIPLELQRLVHGGQVNLDMEDHQDQEYIKPRL RFKAFSGEGQKLGSLTPEIVSTPSSPEEEDKSILNAVVLIDDSVPTTKIQIRLADGSRLIQRFNSTHRIL

DVRNFIVQSRPEFAALDFILVTSFPNKELTDESLTLLEADILNTVLLQQLK

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK
Predicted MW: 36.9 kDa

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 001071087

**Locus ID:** 137886






**UniProt ID:** Q14CS0 5088 RefSeq Size: Cytogenetics: 8q12.1 993 RefSeq ORF: Synonyms: p37

**Summary:** Adapter protein required for Golgi and endoplasmic reticulum biogenesis

> (PubMed:17141156). Involved in Golgi and endoplasmic reticulum maintenance during interphase and in their reassembly at the end of mitosis (PubMed:17141156). The complex formed with VCP has membrane fusion activity; membrane fusion activity requires USO1-GOLGA2 tethering and BET1L (PubMed:17141156). VCPIP1 is also required, but not its deubiquitinating activity (PubMed:17141156). Together with NSFL1C/p47, regulates the centrosomal levels of kinase AURKA/Aurora A during mitotic progression by promoting AURKA removal from centrosomes in prophase (PubMed:23649807). Also, regulates spindle

orientation during mitosis (PubMed:23649807).[UniProtKB/Swiss-Prot Function]

## **Product images:**



Coomassie blue staining of purified UBXN2B protein (Cat# TP319137). The protein was produced from HEK293T cells transfected with UBXN2B cDNA clone (Cat# [RC219137]) using MegaTran 2.0 (Cat# [TT210002]).