

Product datasheet for TP319253

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

HCST (NM_001007469) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human hematopoietic cell signal transducer (HCST), transcript variant

2, 20 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC219253 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MIHLGHILFLLLPVAAAQTTPGERSSLPAFYPGTSGSCSGCGSLSLPLLAGLVAADAVASLLIVGAVFL

CARPRRSPAQEDGKVYINMPGRG

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 7.3 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeg: NP 001007470

 Locus ID:
 10870

 UniProt ID:
 Q9UBK5

RefSeq Size: 521





Cytogenetics: 19q13.12

RefSeq ORF: 279

Synonyms: DAP10; KAP10; PIK3AP


Summary: This gene encodes a transmembrane signaling adaptor that contains a YxxM motif in its

> cytoplasmic domain. The encoded protein may form part of the immune recognition receptor complex with the C-type lectin-like receptor NKG2D. As part of this receptor complex, this protein may activate phosphatidylinositol 3-kinase dependent signaling pathways through its intracytoplasmic YxxM motif. This receptor complex may have a role in cell survival and proliferation by activation of NK and T cell responses. Alternative splicing results in two

transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome, Transmembrane **Protein Pathways:** Natural killer cell mediated cytotoxicity

Product images:



Coomassie blue staining of purified HCST protein (Cat# TP319253). The protein was produced from HEK293T cells transfected with HCST cDNA clone (Cat# [RC219253]) using MegaTran 2.0 (Cat# [TT210002]).