

Product datasheet for **TP319336L**

IDUA (NM_000203) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins
Description: Purified recombinant protein of Homo sapiens iduronidase, alpha-L- (IDUA), 1 mg
Species: Human
Expression Host: HEK293T
Expression cDNA >RC219336 representing NM_000203
Clone or AA Sequence: **Red**=Cloning site **Green**=Tags(s)

MRPLRPRAALLALLASLLAAPPVAPAEAPHLVHVDAARALWPLRRFWRSTGFCPPLPHSQADQYVLSWDQ
QLNLAYVGVAVPHRGIKQVRTHWLELVTTTRGSTGRGLSYNFTHLDGYLDLLRENQLLPGFELMGSASGHF
TDFEDKQQVFEWKDLVSSLARRYIGRYGLAHVSKWNFETWNEPDHDFDNVSMQMGFLNYYDACSEGLR
AASPALRLGGPGDSFHTPPRSPLSWGLLRHCHDGTNFFTGEAGVRLDYISLHRKGARSSISILEQEKVVA
QQIRQLFPKFADTPIYNDEADPLVGWVSLPQWRADVTYAAAMVVKVIAQHQNLLANTTSAPFYALLSNDN
AFLSYHPHPFAQRTLARFQVNNTRPPHVQLLRKPVLTAMGLLALLDEEQLWAEVSQAGTVLDSNHTVGV
LASAHRPQGPADAWRAAVLIASDDTRAHPNRSVAVTLRLRGVPPGGLVYVTRYLDNGLCSPDGEWRRR
GRPVFPTAEQFRMRAAEDPVAAPRPLPAGGRLLRPLALRPLSLLLHVHVCARPEKPPGQVTRLRALPLT
QGQLVLVWSDEHVGSKCLWYIEIQFSQDGKAYTPVSRKPSTFNLVFPSPDTGAVSGSYRVRALDYWARPG
PFSDVPYLEVPVPRGPPSPGNP

SGPTRRRLEQKLISEEDLAANDILDYKDDDDKV


Tag: C-Myc/DDK
Predicted MW: 70 kDa
Concentration: >0.05 µg/µL as determined by microplate BCA method
Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Bioactivity: Not mentioned (PMID: [29671261](#))
Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.



[View online »](#)

Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_000194
Locus ID:	3425
UniProt ID:	P35475
RefSeq Size:	2197
Cytogenetics:	4p16.3
RefSeq ORF:	1959
Synonyms:	IDA; MPS1; MPSI
Summary:	This gene encodes an enzyme that hydrolyzes the terminal alpha-L-iduronic acid residues of two glycosaminoglycans, dermatan sulfate and heparan sulfate. This hydrolysis is required for the lysosomal degradation of these glycosaminoglycans. Mutations in this gene that result in enzymatic deficiency lead to the autosomal recessive disease mucopolysaccharidosis type I (MPS I). [provided by RefSeq, Jul 2008]
Protein Families:	Druggable Genome
Protein Pathways:	Glycosaminoglycan degradation, Lysosome, Metabolic pathways

Product images:



Coomassie blue staining of purified IDUA protein (Cat# [TP319336]). The protein was produced from HEK293T cells transfected with IDUA cDNA clone (Cat# [RC219336]) using MegaTran 2.0 (Cat# [TT210002]).