

Product datasheet for TP319713L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

KCTD1 (NM_198991) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human potassium channel tetramerisation domain containing 1

(KCTD1), transcript variant 2, 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC219713 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MSRPLITRSPASPLNNQGIPTPAQLTKSNAPVHIDVGGHMYTSSLATLTKYPESRIGRLFDGTEPIVLDS LKQHYFIDRDGQMFRYILNFLRTSKLLIPDDFKDYTLLYEEAKYFQLQPMLLEMERWKQDRETGRFSRPC ECLVVRVAPDLGERITLSGDKSLIEEVFPEIGDVMCNSVNAGWNHDSTHVIRFPLNGYCHLNSVQVLERL

QQRGFEIVGSCGGGVDSSQFSEYVLRRELRRTPRVPSVIRIKQEPLD

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 29.2 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 945342

Locus ID: 284252



KCTD1 (NM_198991) Human Recombinant Protein - TP319713L

UniProt ID: <u>Q719H9</u>, <u>A0A024RC45</u>

RefSeq Size: 1754

Cytogenetics: 18q11.2

RefSeq ORF: 771

Synonyms: C18orf5


Summary: This gene encodes a protein containing a BTB (Broad-complex, tramtrack and bric a brac),

also known as a POZ (POxvirus and zinc finger) protein-protein interaction domain. The encoded protein negatively regulates the AP-2 family of transcription factors and the Wnt signaling pathway. A mechanism for the modulation of Wnt signaling has been proposed in which the encoded protein enhances ubiquitination and degradation of the beta-catenin protein. Mutations in this gene have been identified in Scalp-ear-nipple (SEN) syndrome.

[provided by RefSeq, May 2017]

Protein Families: Ion Channels: Other

Product images:



Coomassie blue staining of purified KCTD1 protein (Cat# [TP319713]). The protein was produced from HEK293T cells transfected with KCTD1 cDNA clone (Cat# [RC219713]) using MegaTran 2.0 (Cat# [TT210002]).