

Product datasheet for TP319855

TGF beta Receptor II (TGFB2) (NM_003242) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human transforming growth factor, beta receptor II (70/80kDa) (TGFB2), transcript variant 2, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC219855 representing NM_003242 Red=Cloning site Green=Tags(s)

MGRGLLRGLWPLHIVLWTRIASTIPPHVQKSVNNDMIVTDNNGAVKFPQLCKFCDVRFSTCDNQKSCMSN
CSITSICEKPEVCVAVWRKNDENITLETVCHDPKLPYHDFILEDAAAPKCIMKEKKKPGETFFMCSCSS
DECNDNIIFSEYNTSNPDLLLIVFQVTGISLLPPLGVAISVIIIIFYCYRVNRQQKLSSTWETGKTRKLM
EFSEHCAIILEDSDISSTCANNINHNTPELLIEDTLVVGKGRFAEVYKAKLKQNTSEQFETVAVKIFP
YEEYASWKTEKDIFSDINLKHENILQFLTAERKTELKQYWLITAFHAKGNLQEYLTRHVISWEDLRKL
GSSLARGIAHLHSDHTPCGRPKMPIVHRDLKSSNILVKNDLTCCCLDFGLSLRLDPTLSVDDLANSGQVG
TARYMAPEVLESRMNLENVESFKQTDVYSMALVLWEMTSRCNAVGEVKDYEPFGSKVREHPCVESMKDN
VLRDRGRPEIPSWLNLHQIQMVCETLTECWDHDPPEARLTAQCVAERFSELEHLDRLSGRSCSEEKIPED
GSLNNTK

SGPTRRRLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	62 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_003233
Locus ID:	7048
UniProt ID:	P37173 , A3QNQ0
RefSeq Size:	4639
Cytogenetics:	3p24.1
RefSeq ORF:	1701
Synonyms:	AAT3; FAA3; LDS1B; LDS2; LDS2B; MFS2; RIIC; TAAD2; TBR-ii; TBRli; TGFbeta-RII; TGFR-2
Summary:	The protein encoded by this gene is a transmembrane protein that has a protein kinase domain, forms a heterodimeric complex with TGF-beta receptor type-1, and binds TGF-beta. This receptor/ligand complex phosphorylates proteins, which then enter the nucleus and regulate the transcription of genes related to cell proliferation, cell cycle arrest, wound healing, immunosuppression, and tumorigenesis. Mutations in this gene have been associated with Marfan Syndrome, Loeys-Deitz Aortic Aneurysm Syndrome, and the development of various types of tumors. Alternatively spliced transcript variants encoding different isoforms have been characterized. [provided by RefSeq, Aug 2017]
Protein Families:	Druggable Genome, Protein Kinase, Transmembrane
Protein Pathways:	Adherens junction, Chronic myeloid leukemia, Colorectal cancer, Cytokine-cytokine receptor interaction, Endocytosis, MAPK signaling pathway, Pancreatic cancer, Pathways in cancer, TGF-beta signaling pathway

Product images:



Coomassie blue staining of purified TGFB2 protein (Cat# TP319855). The protein was produced from HEK293T cells transfected with TGFB2 cDNA clone (Cat# [RC219855]) using MegaTran 2.0 (Cat# [TT210002]).