OriGene Technologies, Inc.
9620 Medical Center Drive, Ste 200
Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Product datasheet for TP319913L

DGKE (NM_003647) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human diacylglycerol kinase, epsilon 64kDa (DGKE), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC219913 representing NM_003647 Red=Cloning site Green=Tags(s) MEAERRPAPGSPSEGLFADGHLILWTLCSVLLPVFITFWCSLQRSRRQLHRRDIFRKSKHGWRDTDLFSQ PTYCCVCAQHILQGAFCDCCGLRVDEGCLRKADKRFQCKEIMLKNDTKVLDAMPHHWIRGNVPLCSYCMV CKQQCGCQPKLCDYRCIWCQKTVHDECMKNSLKNEKCDFGEFKNLIIPPSYLTSINQMRKDKKTDYEVLA SKLGKQWTPLIILANSRSGTNMGEGLLGEFRILLNPVQVFDVTKTPPIKALQLCTLLPYYSARVLVCGGD GTVGWVLDAVDDMKIKGQEKYIPQVAVLPLGTGNDLSNTLGWGTGYAGEIPVAQVLRNVMEADGIKLDRW KVQVTNKGYYNLRKPKEFTMNNYFSVGPDALMALNFHAHREKAPSLFSSRILNKAVYLFYGTKDCLVQEC KDLNKKVELELDGERVALPSLEGIIVLNIGYWGGGCRLWEGMGDETYPLARHDDGLLEVVGVYGSFHCAQ IQVKLANPFRIGQAHTVRLILKCSMMPMQVDGEPWAQGPCTVTITHKTHAMMLYFSGEQTDDDISSTSDQ EDIKATE TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	63.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.

View online »

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_003638
Locus ID:	8526
UniProt ID:	P52429, A1L4Q0
RefSeq Size:	2562
Cytogenetics:	17q22
RefSeq ORF:	1701
Synonyms:	AHUS7; DAGK5; DAGK6; DGK; NPHS7
Summary:	Diacylglycerol kinases are thought to be involved mainly in the regeneration of phosphatidylinositol (PI) from diacylglycerol in the PI-cycle during cell signal transduction. When expressed in mammalian cells, DGK-epsilon shows specificity for arachidonyl-containing diacylglycerol. DGK-epsilon is expressed predominantly in testis. [provided by RefSeq, Jul 2008]
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	Glycerolipid metabolism, Glycerophospholipid metabolism, Metabolic pathways, Phosphatidylinositol signaling system

Product images:



Coomassie blue staining of purified DGKE protein (Cat# [TP319913]). The protein was produced from HEK293T cells transfected with DGKE cDNA clone (Cat# [RC219913]) using MegaTran 2.0 (Cat# [TT210002]).