

Product datasheet for **TP320309**

PLA2G2F (NM_022819) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human phospholipase A2, group IIF (PLA2G2F), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC220309 representing NM_022819 Red =Cloning site Green =Tags(s) MADGAKANPKGFKKKVLDRCFSGWRGPRFGASCPSRTRSRLGMKKFFTVAILAGSVLSTAHGSLNLKA MVEAVTGRSAILSFVGYGCYCYGLGRRGQPKDEVDWCCHAHDCCYQELFDQGCHPYVDHYDHTIENTEI V CSDLNKTECDKQTCMCDKNMVLCLMNQTYREEYRGFLNVYCGPTPNCSIYEPPEEVTCSHQSPAPPA P P TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	23.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_073730
Locus ID:	64600



[View online »](#)

UniProt ID:	<u>Q9BZM2</u>
RefSeq Size:	2737
Cytogenetics:	1p36.12
RefSeq ORF:	633
Synonyms:	GIIFsPLA2; sPLA2-IIF
Summary:	<p>May play a role in lipid mediator production in inflammatory conditions, by providing arachidonic acid to downstream cyclooxygenases and lipoxygenases (By similarity). Phospholipase A2, which catalyzes the calcium-dependent hydrolysis of the 2-acyl groups in 3-sn-phosphoglycerides (PubMed:11112443). Hydrolyzes phosphatidylethanolamine more efficiently than phosphatidylcholine, with only a modest preference for arachidonic acid versus linoelic acid at the sn-2 position. Comparable activity toward 1-palmitoyl-2-oleoyl-phosphatidylserine vesicles to that toward 1-palmitoyl-2-oleoyl-phosphatidylglycerol (By similarity). Hydrolyzes phosphatidylglycerol versus phosphatidylcholine with a 15-fold preference (PubMed:11112443).[UniProtKB/Swiss-Prot Function]</p>
Protein Families:	Druggable Genome, Secreted Protein
Protein Pathways:	alpha-Linolenic acid metabolism, Arachidonic acid metabolism, Ether lipid metabolism, Fc epsilon RI signaling pathway, Glycerophospholipid metabolism, GnRH signaling pathway, Linoleic acid metabolism, Long-term depression, MAPK signaling pathway, Metabolic pathways, Vascular smooth muscle contraction, VEGF signaling pathway