

Product datasheet for **TP320400**

LIM kinase 2 (LIMK2) (NM_016733) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human LIM domain kinase 2 (LIMK2), transcript variant 2b, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC220400 representing NM_016733 Red=Cloning site Green=Tags(s)

MGSYLSVPAYFTSRDLFRCSECQDSL TNWYYEKDGKLYCPKDYWGKFGFCHGCSLLMTGPFMVAGEFKY
HPECFACMSCKVIIEDGDAYALVQHATLYCGKCHNEVLAPMFERLSTESVQEQLPYSVTLISMPATTEG
RRGFSVSVESACS NYATTVQVKEVNRMHISPNNRNAIHPGDRILEINGTPVRTLRVEEVEDAISQTSQTL
QLLIEHDPVSQRDLQRLRLRLAPHMQNAGHPHALSTLDTKENLEGLTRRRSLRRSNSISKSPGPSSPKE
PLLFSRDISRSESLRCSYSYQQIFRPCDLIHGEVLGKGGFFGQAIKVTHKATGKVMVMKELIRCDEETQK
TFLTEVKVMRSLDHPNVLKFIGVLYKDKKLNLLTEYIEGGTLKDFLRSMDFPWPQQKVRFAKGIASGMAY
LHSMCIHRDLNSHNCLIKLDKTVVADFGLSRLIVEERKRAPMEKATTKRTRLNDRKKRYTVVGNPY
WMAPEMLNGKSYDETVDIFSFGIVLCEIIGQVYADPDCLPRTLDFGLNVKLFWEKFPVPTDCPPAFFPLAA
ICCRLEPESRPAFSKLEDSFEALSPLYLGELGIPLPAELEELDHTVSMQYGLTRDSP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	69.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_057952](#)

Locus ID: 3985

UniProt ID: [P53671](#), [A0A024R1M2](#)

RefSeq Size: 3848

Cytogenetics: 22q12.2


RefSeq ORF: 1851

Summary: There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. The protein encoded by this gene is phosphorylated and activated by ROCK, a downstream effector of Rho, and the encoded protein, in turn, phosphorylates cofilin, inhibiting its actin-depolymerizing activity. It is thought that this pathway contributes to Rho-induced reorganization of the actin cytoskeleton. At least three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: Axon guidance, Fc gamma R-mediated phagocytosis, Regulation of actin cytoskeleton

Product images:



Coomassie blue staining of purified LIMK2 protein (Cat# TP320400). The protein was produced from HEK293T cells transfected with LIMK2 cDNA clone (Cat# [RC220400]) using MegaTran 2.0 (Cat# [TT210002]).