

Product datasheet for **TP320496L**

BMT2 (NM_152556) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human chromosome 7 open reading frame 60 (C7orf60), 1 mg

Species: Human

Expression Host: HEK293T

**Expression cDNA Clone
or AA Sequence:** >RC220496 protein sequence
Red=Cloning site **Green**=Tags(s)

MEPGAGGRNTARAQRAGSPNTPPPREQERKLEQEKLSGVVKSVHRRRLRKKYREVGDFDKIWREHCEDEET
LCEYAVAMKNLADNHWAKTCEGEGRIEWCCSVCREYFQNGGKRKALEKDEKRAVLATKTTPALNMHESQ
LEGHLTNLSFTNPEFITELLQASGKIRLLDVGSCFNPFLKFEFLTVGIDIVPAVESVYKCDFLNLQLQQ
PLQLAQDAIDAFLKQLKNPIDSLPGELFHVVVFSLLSYFSPYQRWICCKKAHELLVLNGLLLIITPDS
SHQNRHAMMMKSWKIAIESLGFKRFKYSKFSMHMLMAFRKISLKTSDLVSRNYPGMLYIPQDFNSIEDE
EYSNPSCYVRSIDIEDEQLAYGFTLPDAPYDSDSGESQASSIPFYELEDPIILLS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 46.1 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_689769](#)

Locus ID: 154743



[View online »](#)

UniProt ID: [Q1RMZ1](#)

RefSeq Size: 3969


Cytogenetics: 7q31.1

RefSeq ORF: 1215

Synonyms: C7orf60; SAMTOR

Summary: S-adenosyl-L-methionine-binding protein that acts as an inhibitor of mTORC1 signaling via interaction with the GATOR1 and KICSTOR complexes (PubMed:29123071). Acts as a sensor of S-adenosyl-L-methionine to signal methionine sufficiency to mTORC1: in presence of methionine, binds S-adenosyl-L-methionine, leading to disrupt interaction with the GATOR1 and KICSTOR complexes and promote mTORC1 signaling (PubMed:29123071). Upon methionine starvation, S-adenosyl-L-methionine levels are reduced, thereby promoting the association with GATOR1 and KICSTOR, leading to inhibit mTORC1 signaling (PubMed:29123071). Probably also acts as a S-adenosyl-L-methionine-dependent methyltransferase (Potential).[UniProtKB/Swiss-Prot Function]

Product images:



Coomassie blue staining of purified BMT2 protein (Cat# [TP320496]). The protein was produced from HEK293T cells transfected with BMT2 cDNA clone (Cat# [RC220496]) using MegaTran 2.0 (Cat# [TT210002]).