

## Product datasheet for TP320503M

### JAK2 (NM\_004972) Human Recombinant Protein

#### Product data:

|                                       |  |
|---------------------------------------|--|
| Product Type:                         | Recombinant Proteins   |
| Description:                          | Recombinant protein of human Janus kinase 2 (a protein tyrosine kinase) (JAK2), 100 µg |
| Species:                              | Human  |
| Expression Host:                      | HEK293T  |
| Expression cDNA Clone or AA Sequence: | >RC220503 representing NM_004972<br>Red=Cloning site Green=Tags(s)                     |

MGMACLTMTMEGTSTSSYQNGDISGNANSMKQIDPVLQVYLYHSLGKSEADYLTFFPSGEYVAEEICIA  
ASKACGITPVYHNMFALMSETERIWYPPNHVHFHIDESTRHNVLYRIRFYFPRWYCSGNSRAYRHGISRGA  
EAPLLDDFVMSYLFAQWRHDFVHGWIQVPVTHETQEECLGMAVLDMMRIAKENDQTPLAIYNSISYKTFL  
PKCIRAKIQDYHILTRKRIRYRFRRIQFSQCKATARNLKLKYLINLETLSAFYTEKFEVKEPGSGPS  
GEEIFATIIITGNGGIQWSRGKHKESETLTEQDLQLYCDFPNIIDVSIKQANQEGSNESRVVTIHKQDGK  
NLEIELSSLREALSFVSLIDGYRRLTADAHHYLCKEVAPPVLENIQSNCHGPISMDFAISLKKAGNQT  
GLYVLRCSPKDFNKYFLTFEVERENVIEYKHCLITKNENEEYNLSGTTKKNFSSLDLLNCYQMETVRSND  
IIFQFTKCCPPKPKDKSNLLVFRNTGVSVDVPTSPQLRPTHMNQMVFHKIRNEDLIFNESLGQGTFTKIF  
KGVRRVVDYDGLHETEVLLKVLDAHRNYSSEFFEAASMMSKLSHKHLVLYGVCVCGDENILVQEFVK  
FGSLDTYLLKKNKNCINILWKLEVAQQLAWAMHFLEENTLIHGNVCAKNILLIREDRKTGNPPFIKLSDP  
GISITVLPKDILQERIPWVPECIENPKNLNLATDKWSFGTTLWEICSGGDKPLSALDSQRKLQFYEDRH  
QLPAPKWAELANLINCMYEPDFRPSFRAIIRDLNLSLFTPDYELLTENDMLPNMRIGALGFSGAFEDRD  
PTQFEERHLKFLQQLGKGNFGSVEMCRYDPLQDNTGEVAVKQLQHSTEEHLRDFEREIEILKSLQHDNI  
VKYKGVCSAGRRNLKLIMEYLPYGLRDYLLQKHKERIDHIKLLQYTSQICKGMEYLGTKRYIHRDLATR  
NILVENENRVKIGDFGLTKVLPQDKEYYKVKEPGESPIFWYAPESLTSKFSVSDVWVSGVWLYELFTY  
IEKSKSPPAEFMRMIGNDKQGMIVFHLLIELLKNNGRLLPRPDGCPDEIYMIMTECWNNNVNQRPSFRDLA  
LRVDQIRDNMAG


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

|                |   |
|----------------|---|
| Tag:           | C-Myc/DDK   |
| Predicted MW:  | 130.5 kDa   |
| Concentration: | >0.05 µg/µL as determined by microplate BCA method          |
| Purity:        | > 80% as determined by SDS-PAGE and Coomassie blue staining |
| Buffer:        | 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol        |



[View online »](#)

|                          |   |
|--------------------------|---|
| <b>Preparation:</b>      | Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.  |
| <b>Note:</b>             | For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.  |
| <b>Storage:</b>          | Store at -80°C.   |
| <b>Stability:</b>        | Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.   |
| <b>RefSeq:</b>           | <a href="#">NP_004963</a>   |
| <b>Locus ID:</b>         | 3717  |
| <b>UniProt ID:</b>       | <a href="#">O60674</a>  |
| <b>RefSeq Size:</b>      | 5097  |
| <b>Cytogenetics:</b>     | 9p24.1  |
| <b>RefSeq ORF:</b>       | 3396  |
| <b>Synonyms:</b>         | JTK10   |
| <b>Summary:</b>          | <p>This gene encodes a non-receptor tyrosine kinase that plays a central role in cytokine and growth factor signalling. The primary isoform of this protein has an N-terminal FERM domain that is required for erythropoietin receptor association, an SH2 domain that binds STAT transcription factors, a pseudokinase domain and a C-terminal tyrosine kinase domain. Cytokine binding induces autophosphorylation and activation of this kinase. This kinase then recruits and phosphorylates signal transducer and activator of transcription (STAT) proteins. Growth factors like TGF-beta 1 also induce phosphorylation and activation of this kinase and translocation of downstream STAT proteins to the nucleus where they influence gene transcription. Mutations in this gene are associated with numerous inflammatory diseases and malignancies. This gene is a downstream target of the pleiotropic cytokine IL6 that is produced by B cells, T cells, dendritic cells and macrophages to produce an immune response or inflammation. Disregulation of the IL6/JAK2/STAT3 signalling pathways produces increased cellular proliferation and myeloproliferative neoplasms of hematopoietic stem cells. A nonsynonymous mutation in the pseudokinase domain of this gene disrupts the domains inhibitory effect and results in constitutive tyrosine phosphorylation activity and hypersensitivity to cytokine signalling. This gene and the IL6/JAK2/STAT3 signalling pathway is a therapeutic target for the treatment of excessive inflammatory responses to viral infections. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Jul 2020]</p> |
| <b>Protein Families:</b> | Druggable Genome, Protein Kinase  |
| <b>Protein Pathways:</b> | Adipocytokine signaling pathway, Chemokine signaling pathway, Jak-STAT signaling pathway  |

**Product images:**

Coomassie blue staining of purified JAK2 protein (Cat# [TP320503]). The protein was produced from HEK293T cells transfected with JAK2 cDNA clone (Cat# [RC220503]) using MegaTran 2.0 (Cat# [TT210002]).