

## Product datasheet for **TP320585M**

### **BIN1 (NM\_139344) Human Recombinant Protein**

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human bridging integrator 1 (BIN1), transcript variant 2, 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC220585 representing NM_139344 <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MAEMGSKGVTAGKIASNVQKKLTRAQEKVLQKLGKADETKDEQFEQCVQNFNKQLTEGTRLQKDLRITYLA  
SVKAMHEASKKLNECLQEVYEPDWPGRDEANKIAENNDLLWMDYHQKLVQDQALLTMDTYLQGFPDIKSRI  
AKRGRKLVYDSARHHYESLQTAKKKDEAKIAPVSLLEKAAPQWCQGKLAHLVAQTNLLRNQAEELI  
KAQKVFEEMNVDLQEELPSLWNSRVGFYVNTFQSIAGLEENFHKEMSKLNQNLNDVLVGLEKQHGSNTFT  
VKAQPSDNAPAKGNKSPSPDGSPAATPEIRVNHEPEPAGGATPGATLPKSPSQFEAPGPFSEQASLLDL  
DFDPLPPVTSPVKAPTPSGQSIPWDLWEPTESPAGSLPSGEPSSAAEGTFAVSWPSQTAEPGPAQPAEASE  
VAGGTQPAAGAQPGETAASEAASSSLPAVVVETFPATVNGTVEGGSGAGRLDLPFGFMFKVQAQHDYTA  
TDTDELQKAGDWLVVIPFQNPEEQDEGWLMGVKESDWNQHKELEKCRGVFPENFTEPVP

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**

Tag:	C-Myc/DDK
Predicted MW:	59.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.




[View online >](#)

RefSeq:	<a href="#">NP_647594</a>
Locus ID:	274
UniProt ID:	<a href="#">O00499</a> , <a href="#">A0A024RAF6</a>
RefSeq Size:	2508
Cytogenetics:	2q14.3
RefSeq ORF:	1650
Synonyms:	AMPH2; AMPHL; CNM2; SH3P9

**Summary:** This gene encodes several isoforms of a nucleocytoplasmic adaptor protein, one of which was initially identified as a MYC-interacting protein with features of a tumor suppressor. Isoforms that are expressed in the central nervous system may be involved in synaptic vesicle endocytosis and may interact with dynamin, synaptojanin, endophilin, and clathrin. Isoforms that are expressed in muscle and ubiquitously expressed isoforms localize to the cytoplasm and nucleus and activate a caspase-independent apoptotic process. Studies in mouse suggest that this gene plays an important role in cardiac muscle development. Alternate splicing of the gene results in several transcript variants encoding different isoforms. Aberrant splice variants expressed in tumor cell lines have also been described. [provided by RefSeq, Mar 2016]

### Product images:



Coomassie blue staining of purified BIN1 protein (Cat# [TP320585]). The protein was produced from HEK293T cells transfected with BIN1 cDNA clone (Cat# [RC220585]) using MegaTran 2.0 (Cat# [TT210002]).